



## EFFECT OF NITROGEN AND VAM LEVELS ON HERBAGE AND OIL YIELD OF PATCHOULI (*Pogostemon patchouli* Petle.)

<sup>a</sup>Sumathi. M., <sup>b</sup>Shashekala, S.G., <sup>c</sup>Shankaraiah. N., <sup>d</sup>Ravi Kumar. P. & <sup>e</sup>Kavitha, V.

<sup>a</sup>M.S.Swaminathan Research Foundation, Chennai

<sup>b</sup>IFFCO Kisan Sanchar limited, Bangalore

<sup>c</sup>Indian Institute of Horticultural Research, Hessaraghatta, Bangalore

<sup>d</sup>Department of Animal Husbandry and veterinary services, Government of Karnataka)

<sup>e</sup>Department of Agricultural Economics, Tamil Nadu Agricultural University, Tamil Nadu as fifth author.

### ABSTRACT

*Pogostemon patchouli* Pellet. Is an important aromatic plant, which gains commercial importance in recent years. The increasing demand for the essential oil both for the perfumery industry and pharmaceutical industry stressed the need for systematic investigation for increasing the herbage and oil yield. An experiment was therefore carried out in Horticultural College and Research Institute in Periyakulam, to find the effect of different levels of Nitrogen and VAM levels on herbage and oil yield Patchouli. The study revealed that, effect of nitrogen and VAM had significant effect on yield and yield attributing characters. The plant height was increased with the increased dose of nitrogen and VAM application, the maximum plant height was recorded with 200kg/ha nitrogen and 50kg/ha VAM ( 89.51 cm). Application of 150kg/ha nitrogen and 50kg/ha VAM recorded highest number of laterals ( 20.27). Number of leaves was increased with application of nitrogen up to 15kg/ha and the highest number of leaves (361) was recorded with application of 150kg N/ha along with 50kg VAM/ha. The highest dose of nitrogen and VAM had significant influence on leaf area and leaf area index. Application of 200kg nitrogen with 50kg VAM/ha recorded the maximum leaf area and leaf area index but it was on par with nitrogen at 150kg/ha and VAM at 50kg/ha. Highest herbage yield (13.60 t/ha) and dry weight of herbage (4.07 t/ha) was recorded at 150kg N/ha and 5kg VAM/ha. Essential oil content of the treatments did not varied significantly among the treatments. Essential oil yield in herbage was highest at 150kg nitrogen and 50kg VAM/ha(110.42 kg/ha). Highest mycorrhizal colonization percentage of 90.50% was recorded with the treatment receiving 150kg nitrogen/ha and 50 kg VAM/ha. In the nutshell of the research, application of Nitrogen @150kg/ha and VAM at 50kg/ha enhanced the quantitative and qualitative traits and recorded maximum cost benefit ratio (1: 3.67).

**KEY WORDS:** Farm Yard Manure, Nitrogen, Phosphorus, Potassium, VAM & DAP.

### INTRODUCTION

Patchouli (*Pogostemon patchouli* Pellet.) syn. (*Pogostemon cablin* Benth.) is one of the important aromatic crops which belongs to the family Lamiaceae and its native to Philippines. Shade dried leaves of Patchouli on steam distillation yield essential oil containing about 97 per cent of compounds which have no influence on aroma, out these 40 to 45 per cent belongs to sesquiterpene group and the balance seems to consist of patchouli alcohol. Patchouli oil is one of the most important essential oil of the perfumery industry, as the oil blends well worth other essential oils like vetiver, sandal wood, geranium, lavender, clove oil etc. Hence it is regarded as the best fixative for heavy perfumes imparting strength, character and alluring notes and lasting qualities. Besides the oil is also used as flavor ingredients in the major food products, including alcoholic and nonalcoholic beverages. In Indo – Malayan region, it has been used as insecticide and leach repellent and also to soothe menstrual cramps. There is no synthetic chemical to replace the oil of patchouli which further enhances its value (Ramachandra *et al.*, 2002). Thus, it has a unique position in the market and the oil is

in great demand in perfumery, soap, scent, after shave lotion, detergent, tobacco and incense manufacturing industries. It has been amply shown that patchouli being a tropical plant can be successfully grown under Indian conditions (Bhaskar *et al.*, 1997.) Around 1962, systemic efforts on cultivation were started by Central Institute of Medicinal and Aromatic plants (CIMAP) at regional Centre, Bangalore. Production of Patchouli in India is very negligible (about 100 – 150kg/year), as against the total annual world production of 700 to 800 tones. Presently India is importing over 200 tones of oils from Indonesia, Malaysia and Singapore ( Manjunatha *et al.*, 2002).

To grow any crop, the point of paramount importance is to get maximum output with minimum of inputs. This can be achieved through evolution of scientifically sound and economically feasible methodologies in the management of crops. This includes standardization of nutritional requirements. Some research work in patchouli has already been carried out to find out optimum fertilizer requirements etc. But these findings are to be suitably fine-tuned by further research to suit under Tamil Nadu conditions.

Among the various nutrients, nitrogen, phosphorus, potassium are the three important nutrients that are

frequently short supply in the soil and their application plays a very important role in altering various growth, yield and quality attributes of the plants. However, the modern and intensive agriculture calls for the heavy dependence of fertilizers, and chemicals, which are not only costly but also cause soil and water pollution. Thus, by considering the recent concept of ecofriendly technology, application of bio fertilizers in combination with inorganic fertilizers substitutes the above need in many crops. Thus, keeping the above facts in the view the present investigation was under taken to find out the "Effect of nitrogen and VAM levels on herbage and oil yield of patchouli (*Pogostemon patchouli*) under coconut shade with the following objectives, *i.e.* to fix the optimum dose of nitrogen and VAM levels for patchouli & to maximize herbage and oil yield of patchouli

## MATERIALS AND METHODS

FYM at the rate of 15 tonnes per hectare was incorporated at the time of last ploughing. The recommended dose of 50 kg phosphorus and 50kg potash per hectare were applied in the form of single super phosphate and muriate of potash as a basal dose. In addition uniform dose of biofertilizers viz., Azospirillum and Phosphobacteria, at the rate of 2kg/ha were applied around the root zone of the crops. Tissue culture plants of uniform size variety Johore had planted at uniform spacing of 45 X 45 cm in the plot size of 2.5 X 2.5M which accommodates 30 plants/plot. The experiment was laid out in Factorial Randomized Block Design, replicated twice with five levels of Nitrogen viz., 0(N<sub>1</sub>), 50 (N<sub>2</sub>), 100 (N<sub>3</sub>), 150 (N<sub>4</sub>) and 200 (N<sub>5</sub>) kg/ha, five levels of VAM viz., 0 (V<sub>1</sub>), 12.5 (V<sub>2</sub>), 25 (V<sub>3</sub>), 37.5 (V<sub>4</sub>) and 50 (V<sub>5</sub>) kg/ha. At the time of planting, 1/3<sup>rd</sup> of nitrogen was applied as basal dose and the remaining quantity was applied in two equal split doses after 30<sup>th</sup> and 60<sup>th</sup> days of planning. The entire quantity of VAM (*Glomus fasciculatum*) was applied after 10days of planning by farming small furrows around the root zone of the crop. Periodical hand weeding was carried out to keep the plot free of weeds. Irrigation was given at an interval of 5-6 days depending upon the soil moisture conditions. There was no serious pest and disease incidence observed

throughout the experimental period. However, stray incidences of leaf eating caterpillar ( *Pronomus profusalis*), rhizoctonia wilt and root knot nematode (*Meloidogyne incognita*) were noticed. Matured shoots of 25 to 30 centimeter length were harvested by cutting with sharp secateurs in such a way that the plants retains some of the bottom leaves to ensure better growth at next year. After harvesting herbage was dried for 5 to 6 days by spreading it in a thin layer on hard, dry, cement surface. The leaves were turned periodically to ensure uniform drying and to prevent mould formation. Latter, the dried leaves were steam distilled for oil extraction.

In each treatment, five plants were randomly selected and tagged, biometric observations like plant height, number of laterals, plant spread, leaf area and leaf area index, stalk length, herbage yield and oil content was taken that plants after 180 days of planting. Mycorrhizal parameters like per cent root colonization also observed. Collected data were analyzed adopting the procedure described by panes and Sukhatme (1957), Correlation Coefficients were worked out as suggested by Snedecor and Cochran (1967).

## RESULTS AND DISCUSSION

### Plant height

Nitrogen levels significantly improved plant height and there is a linear relationship was observed in plant height and the levels of nitrogen application. This may be due to the effect of nitrogen in promoting the vegetative growth by the enhanced cell division and the greater synthesis of chlorophyll, protein and amino acids. Irulappan and Ponnusami (1982) and Saha et al (1992) have obtained similar results in Patchouli. The highest value of plant height was recorded with 50kg VAM/ha, this may be due to the effective utilization of VAM that enhanced availability of nutrients along with the production of some growth promoting substances, which might have caused cell elongation and multiplication. Same findings have reported in many horticultural crops like palmarosa (Gupta and Janardhanan, 1991) and African marigold (Rajadurai *et al* 2000).

TABLE -I

|                |       | V <sub>2</sub> | V <sub>3</sub> | V <sub>4</sub> | V <sub>5</sub> | Mean  |
|----------------|-------|----------------|----------------|----------------|----------------|-------|
| N <sub>1</sub> | 35.37 | 39.55          | 44.26          | 49.53          | 50.27          | 43.79 |
| N <sub>2</sub> | 55.38 | 59.46          | 64.28          | 69.33          | 73.30          | 64.35 |
| N <sub>3</sub> | 61.30 | 67.17          | 70.25          | 74.77          | 78.55          | 70.40 |
| N <sub>4</sub> | 68.37 | 71.58          | 75.41          | 79.64          | 85.26          | 76.05 |
| N <sub>5</sub> | 72.54 | 76.96          | 80.46          | 84.15          | 89.51          | 80.72 |
| Mean           | 58.59 | 62.94          | 66.93          | 71.48          | 75.38          | 67.06 |

The interaction between Nitrogen and VAM had a significant effect on plant height and the maximum plant height was observed with 200kg/ha of nitrogen coupled with 50kg VAM/ha. This findings was in agreement with the findings of Shivalingappa (1998) in tuberose.

### Number of laterals per plant

Application of nitrogen at 150kg/ha recorded the maximum numbers of laterals per plant. This may be due

to enhanced vegetative growth because of increased meristamatic activity and increased supply of photosynthates. Arul Arasu and Sambandamurthi (1999) reported similar results in Ocimum.

VAM at different levels significantly increased the number of laterals per plant, the maximum laterals were observed with application of 50kg VAM/ha. Similar results were reported by Manjunatha *et al* (2002) in patchouli. In the treatment that received 150g nitrogen/ha and 50kg

VAM/ha recorded maximum number of laterals followed by 200kg Nitrogen and 50kg VAM/ha.

**TABLE -II**

|                |       | V <sub>2</sub> | V <sub>3</sub> | V <sub>4</sub> | V <sub>5</sub> | Mean  |
|----------------|-------|----------------|----------------|----------------|----------------|-------|
| N <sub>1</sub> | 5.05  | 5.87           | 6.23           | 7.01           | 7.47           | 6.32  |
| N <sub>2</sub> | 8.11  | 9.67           | 11.26          | 13.39          | 14.35          | 11.35 |
| N <sub>3</sub> | 10.36 | 12.06          | 14.23          | 16.12          | 17.10          | 13.97 |
| N <sub>4</sub> | 12.31 | 14.27          | 16.12          | 18.04          | 20.27          | 16.27 |
| N <sub>5</sub> | 12.06 | 14.08          | 16.10          | 17.57          | 19.63          | 15.88 |
| Mean           | 9.57  | 11.19          | 12.85          | 14.42          | 15.76          | 12.72 |

**Number of leaves**

Number of leaves per plant increased with corresponding increase in dose of nitrogen up to 150kg/ha, any further increase in the dose of Nitrogen above this slightly reduces leaf yield. This is because of shedding up of lower leaves due to increased nitrogen application. These results are in accordance with the results of Saha *et al.* (1992) in Patchouli and Chauhan *et al.*(2000) in Palmarosa. VAM application at different levels had a significant influence on number of leaves per plant. Similar results were

reported by Rajadurai *et al* (2000) in african marigold. Present study also indicated that the maximum number of leaves per plant (361.00) was recorded with combined application of 150kg nitrogen/ha along with 50 kg VAM/ha. This may be due to better growing conditions that prevailed in the vicinity of root zone due to the application of VAM that helps the plants to absorb more nutrients from the soil Manjunatha *et al.* (2002) in patchouli.

**TABLE –III**

|                |        | V <sub>2</sub> | V <sub>3</sub> | V <sub>4</sub> | V <sub>5</sub> | Mean   |
|----------------|--------|----------------|----------------|----------------|----------------|--------|
| N <sub>1</sub> | 130.50 | 140.50         | 161.00         | 180.50         | 200.50         | 162.60 |
| N <sub>2</sub> | 230.50 | 241.00         | 261.00         | 281.00         | 305.50         | 263.80 |
| N <sub>3</sub> | 255.50 | 260.50         | 280.50         | 301.00         | 320.50         | 283.60 |
| N <sub>4</sub> | 299.00 | 302.00         | 320.50         | 341.00         | 361.00         | 324.70 |
| N <sub>5</sub> | 284.50 | 291.00         | 311.00         | 331.00         | 352.00         | 313.90 |
| Mean           | 240.00 | 247.00         | 266.80         | 286.90         | 307.90         | 269.72 |

**TABLE –IV**

|                |      | V <sub>2</sub> | V <sub>3</sub> | V <sub>4</sub> | V <sub>5</sub> | Mean |
|----------------|------|----------------|----------------|----------------|----------------|------|
| N <sub>1</sub> | 2.73 | 3.07           | 3.23           | 3.31           | 3.47           | 3.16 |
| N <sub>2</sub> | 3.90 | 3.97           | 4.28           | 4.35           | 4.47           | 4.19 |
| N <sub>3</sub> | 4.23 | 4.48           | 4.64           | 4.83           | 5.01           | 4.64 |
| N <sub>4</sub> | 4.66 | 4.87           | 5.03           | 5.21           | 5.48           | 5.05 |
| N <sub>5</sub> | 4.45 | 4.56           | 4.83           | 5.05           | 5.33           | 4.83 |
| Mean           | 3.99 | 4.19           | 4.40           | 4.55           | 4.75           | 4.38 |

**Stalk length**

Stalk length was significantly influenced by Nitrogen and VAM application.

Nitrogen levels found to significantly influence the stalk length. Application of 150 kg nitrogen/ha recorded maximum stalk length (5.05cm) followed by nitrogen @ 200kg/ha (4.84 cm). Application of 50kg VAM/ha recorded a maximum stalk length (4.75 cm) followed by 37.5kg/ha (4.55 cm and the shortest stalk length was registered in the control (3.99cm).

In the present study interaction between nitrogen and VAM significantly influenced the stalk length. The maximum stalk length of 5.84 cm was reported in the treatment with 150kg nitrogen application coupled with 50kg VAM/ha (5.33 cm) followed by nitrogen at 200kg/ha and VAM at 50kg/ha (5.33 cm).

**Leaf area and leaf area index**

The present study revealed that a linear increase in leaf area with increase in nitrogen levels up to 200kg/ha. This

might be due to increased auxin activity, Carbohydrates and other organic compounds produced as a result of nitrogen application. Increased nitrogen levels caused on increased in chlorophyll synthesis and there by increased the leaf area in *Ocimum sanctum* (Arul Arasu and Sambandamurthi, 1999). VAM application at 50kg/ha recorded maximum leaf area. Application of VAM might have enhanced the availability of nitrogen, phosphorus, and other nutrients along with production of growth hormones which might have increased the length, and breadth of the leaves leading to increased leaf area. Similar observations were recorded by Rajadurai *et al* in african marigold and Manjunatha *et al* (2002) in patchouli. The interaction between Nitrogen and VAM was found to significantly influence the leaf area at all the treatments. Treatment combination receiving 150kg nitrogen/ha and 50kg VAM/ha recorded a maximum leaf area index of 2.01 which was on par with 200kg Nitrogen and 50kg VAM/ha (2.01)

**Plant spread**

Application of nitrogen 15kg/ha recorded maximum plant spread. This may be due to more number of branches. Similar results were obtained by Kiruthika devi (2002) in ashwagandha . Increased VAM application at 50 kg/ha

increased the plant spread. This can be attributed to the production of more number of branches, which in turn have increased the plant spread (Manjunatha *et al.* 2002. in patchouli).

**TABLE –V**

|                |       | V <sub>2</sub> | V <sub>3</sub> | V <sub>4</sub> | V <sub>5</sub> | Mean  |
|----------------|-------|----------------|----------------|----------------|----------------|-------|
| N <sub>1</sub> | 20.60 | 22.00          | 27.91          | 29.89          | 31.47          | 26.37 |
| N <sub>2</sub> | 29.18 | 30.03          | 34.50          | 39.57          | 40.10          | 34.67 |
| N <sub>3</sub> | 31.87 | 33.12          | 38.87          | 45.31          | 47.63          | 39.36 |
| N <sub>4</sub> | 43.59 | 44.34          | 48.66          | 52.50          | 55.58          | 48.93 |
| N <sub>5</sub> | 42.37 | 43.08          | 46.00          | 50.05          | 54.28          | 47.15 |
| Mean           | 33.52 | 34.51          | 39.18          | 43.46          | 45.81          | 39.29 |

The present study indicated that 150 kh nitrogen with 50 kg VAM/ha recorded the highest spread followed by 200kg nitrogen with 50kg VAM/ha. This may be due to increased availability of nutrient elements by VAM, further VAM fungi are known to increase water uptake also (Christopher *et al.*, 1994).

**Fresh herbage yield per hectare**

The individual and pooled effect of fresh herbage yield (g/plant) as influenced by nitrogen and VAM levels were recorded at time of harvest. It could be observed that both individual and interaction effect found significant. The maximum herbage yield (10.64t/ha) was obtained with application of 150kg N/ha followed by 200kg N/ha (10.02). The higher herbage yield may be due to increased

height of plants with more number of laterals per plants as reported by Sadasakthi (1986) in marjoram.

VAM application had a linear effect on herbage yield. Maximum herbage yield of 9.99 t/ha was obtained with application of 50kg VAM/ha followed by 37.5kg/ha (9.08t/ha). Similar results were reported by Krishna Naik (1998) in java citronella and Ratti and Janardhanan (1996) in palmarosa.

The treatment combination of 150kg Nitrogen with 50 Kg VAM/ha recorded maximum herbage yield (13.60t/ha) followed by application of 200kg nitrogen and 50kg VAM/ha (12.83 t/ha) and the lowest herbage yield of 3.16t/ha was obtained with control. This may be due to the increased uptake of both macro and micro by the application of nitrogen and VAM.

**TABLE –VI**

|                |       | V <sub>2</sub> | V <sub>3</sub> | V <sub>4</sub> | V <sub>5</sub> | Mean  |
|----------------|-------|----------------|----------------|----------------|----------------|-------|
| N <sub>1</sub> | 20.60 | 22.00          | 27.91          | 29.89          | 31.47          | 26.37 |
| N <sub>2</sub> | 29.18 | 30.03          | 34.50          | 39.57          | 40.10          | 34.67 |
| N <sub>3</sub> | 31.87 | 33.12          | 38.87          | 45.31          | 47.63          | 39.36 |
| N <sub>4</sub> | 43.59 | 44.34          | 48.66          | 52.50          | 55.58          | 48.93 |
| N <sub>5</sub> | 42.37 | 43.08          | 46.00          | 50.05          | 54.28          | 47.15 |
| Mean           | 33.52 | 34.51          | 39.18          | 43.46          | 45.81          | 39.29 |

**Dry herbage yield**

Dry herbage yield was calculated after 6 days of shade drying. The interaction between nitrogen and VAM was significant in increasing the herbage yield. Application of 150kg n/ha followed by 50kg VAM/ha recorded maximum

herbage yield (4.07) followed by application of 200kg nitrogen and 50kg VAM (3.61t/ha). Similar results were obtained by Manjunatha *et al.* (2002) in patchouli, Earanna *et al.* (2001) in *Coleus aromaticus*.

**TABLE –VII**

|                |      | V <sub>2</sub> | V <sub>3</sub> | V <sub>4</sub> | V <sub>5</sub> | Mean |
|----------------|------|----------------|----------------|----------------|----------------|------|
| N <sub>1</sub> | 0.95 | 0.97           | 1.08           | 1.11           | 1.15           | 1.05 |
| N <sub>2</sub> | 1.22 | 1.62           | 2.02           | 2.52           | 2.57           | 1.99 |
| N <sub>3</sub> | 1.59 | 1.89           | 2.47           | 2.98           | 3.36           | 2.45 |
| N <sub>4</sub> | 2.38 | 2.64           | 3.15           | 3.58           | 4.07           | 3.16 |
| N <sub>5</sub> | 2.25 | 2.45           | 3.13           | 3.53           | 3.61           | 2.99 |
| Mean           | 1.67 | 1.91           | 2.37           | 2.74           | 2.95           | 2.32 |

**Oil content and essential oil yield**

The variation in oil content was not significant among the treatments, however, it varied from 2.15 to 2.90 per cent among the treatments, indicating that nitrogen and VAM have only marginal effect on oil content in patchouli. This

is in line with the findings of Manjunatha *et al.* (2002) in patchouli.

**Essential oil yield per hectare**

Application of nitrogen at 150kg/ha significantly recorded higher oil yield. This may be due to the influence of

nitrogen in promoting the vegetative growth, which resulted in increased herbage production, consequently, essential oil yield increased to greater extent. The similar results were reported by Venugopal (2006) in Patchouli.

The essential oil yield increased with every successive increase in VAM levels and the maximum yield was obtained with 50 kg VAM/ha, this might be due to

influence of VAM in enhancing nutrient uptake, which resulted in increased herbage yield, consequently the essential oil yield per hectare to a greater extent. This is in line with findings of Manjunatha *et al.*, (2002) in patchouli. The present study also revealed that the application of nitrogen 150kg/ha with VAM 50kg/ha (110.42kg/ha).

TABLE –VIII

|                |       | V <sub>2</sub> | V <sub>3</sub> | V <sub>4</sub> | V <sub>5</sub> | Mean  |
|----------------|-------|----------------|----------------|----------------|----------------|-------|
| N <sub>1</sub> | 23.43 | 25.26          | 26.52          | 27.04          | 28.08          | 26.06 |
| N <sub>2</sub> | 31.09 | 42.92          | 52.65          | 62.01          | 63.80          | 50.49 |
| N <sub>3</sub> | 37.90 | 47.13          | 62.18          | 77.86          | 82.98          | 61.61 |
| N <sub>4</sub> | 62.84 | 65.33          | 78.17          | 90.00          | 110.42         | 81.35 |
| N <sub>5</sub> | 56.68 | 58.25          | 72.07          | 81.17          | 93.98          | 72.43 |
| Mean           | 42.38 | 47.77          | 58.31          | 67.61          | 75.85          | 58.38 |

TABLE –IX

|                |      | V <sub>2</sub> | V <sub>3</sub> | V <sub>4</sub> | V <sub>5</sub> | Mean  |
|----------------|------|----------------|----------------|----------------|----------------|-------|
| N <sub>1</sub> | 2.50 | 30.50          | 40.50          | 65.50          | 72.50          | 42.30 |
| N <sub>2</sub> | 4.50 | 60.50          | 68.50          | 70.50          | 75.50          | 55.90 |
| N <sub>3</sub> | 5.50 | 65.50          | 70.00          | 76.50          | 80.50          | 59.60 |
| N <sub>4</sub> | 6.50 | 76.50          | 83.32          | 86.50          | 90.50          | 68.66 |
| N <sub>5</sub> | 6.00 | 72.50          | 80.50          | 85.18          | 88.00          | 66.43 |
| Mean           | 5.00 | 61.10          | 68.56          | 76.83          | 81.40          | 58.57 |

#### Per cent root colonization of VAM

Application of VAM at different levels had significant effect on root colonization.

Nitrogen application at 150kg/ha recorded the higher colonization percent. Higher level of nitrogen triggered the vegetative growth, which in turn increases the colonization percentage. Similar results were obtained by Rajadurai *et al.* (2000) in African marigold. Plants receiving 50kg VAM/ha were found to have higher percentage of root colonization (81.40) followed by VAM at 37.50kg/ha (76.83). whereas the plants in control plot recorded the least colonization of 5.00%. In general, treatments with VAM produced better root colonization, this suggests that the native fungus is not more efficient than the inoculated VAM fungus. Thus, inoculation of plants with fungus helps in better root colonization thereby improving the growth and yield of the crops. This is in agreement with the findings of Sreeramulu *et al.*, (1996) in amaranthus and methi, Madhaiyan *et al.* (2000) in marigold.

Interaction between nitrogen and VAM were found to be statistically significant. Application of nitrogen at 150 kg/ha with VAM 50kg/ha recorded the highest colonization percentage of 90.50 followed by 200kg nitrogen with 50kg VAM/ha (88.00). the lowest colonization percentage was recorded in the control.

#### REFERENCES

Ajimodhin, I., Vasundhara, M., Radhakrishna, D., Biradar, S.L. & Rao, G.G.E. (2005) Integrated nutrient management studies in sweet basil (*Ocimum basilicum* L.) *Indian Perfumer*, 49: 95-101.

Ali, M. S., Yazdani, D., Naghdi Badi, H., Ahwazi, M. & Nazari, F. (1999) Effect of nitrogen and phosphorus fertilizer levels and harvesting schedule on dry matter and oil yield in Peppermint (*Mentha piperita*). *Journal of Medicinal and Aromatic Plant Sciences*, 21 :927-930.

Angadi, S. & Vasanthakumar, T. (1995) Patchouli. In *Advances in Horticulture - Medicinal and Aromatic Plants*, Ed. Chadha K.L. and Gupta, R. Malhotra Publishing house, New Delhi, pp. 751-771.

Arul Arasu, P. and S. Sambandamurthi (1999) Effect of nitrogen and spacing on herbage and oil yield in tulsi (*Ocimum sanctum*). *South Indian Hort.*, 47(1-6): 370-372.

Balyan, S.S. & Sobti, S.N. (1990) Effect of nitrogen, phosphorus and potassium on dry

Balyan, S.S., Sobti, S.N., Pushpangadan, Singh, A. & Atal, C.K. (1982) Cultivation of clocimum at Jammu. In *Cultivation & utilization of aromatic plants*, Ed. Atal, C.K. & Kapur, B.M., Regional Research Laboratory. Jammu-Tawi (India), pp. 481-486.

Bhan, M.K., Kanthi Rekha, Kak, S.N. & Pal, S. (1999) Response of new improved strain RL-931 of Cymbopogon to nitrogen fertilization. *Journal of Medicinal and Aromatic Plant Sciences*, 21: 1027-1029

Bhardwaj, S.D. & Kaushal, A.N. (1990) Nitrogen levels and harvesting management studies on fresh herbage and oil yield in Peppermint cultivars (*Mentha piperita* Linn.) *Indian Perfumer*, 34: 30-41.

- Bhardwaj, S.D., Katoch, P.C. & Kaushal, A.N. (1979) Effect of different levels of nitrogen on herb yield and essential oil content in *Mentha* species. *Indian Journal of Forestry*, 2: 27-30.
- Bhaskar, S. (1995) Growth, herbage and oil yields of patchouli (*Pogostemon patchouli*) as influenced by cultivars and nitrogen fertilization. *Indian Perfumer*, 39: 35-38.
- BHASKAR, S., VASANTHA KUMAR, T. & SRIVASTAVA, H.C., 1997, Influence of growth regulators on production of herbage and oil in patchouli. (*Pogostemon patchouli*). *Indian Perfumer*, 41:98-101.
- Bhaskar, S., Vasanthakumar, T. & Srivastava, H.C. (1998) Growth and yield of scented geranium in relation to nitrogen fertilization. *Journal of Medicinal and Aromatic Plant Sciences*, 20:731-734.
- Bhattacharya, A.K., Rajeshwar Rao, B.R., Kaul, P.N., Singh, K. & Singh, C.P. (1995) Response of rose scented geranium to plant growth regulators. *Indian Perfumer*, 39: 99-101.
- Chalapathi, M. V., Vasundhara, M., Ganghadraeshwar Rao, G., Thimmegowda, S., Prakasa Rao, E.V.S. & Devakumar, N. (2004) Influence of integrated nutrient management on yield of Davana (*Artemisiapallens*). *Indian Perfumer*, 48: 311-316.
- Chattopadhyay, T.K. (1994) Light and temperature relations of fruit crops. In *A Text Book of Pomology*, Ed. Chattopadhyay, T.K., Kalyani publishers, West Bengal, pp. 44-59.
- Chauhan, H.S., Kalra, A., Mengi, N., Rajput, D.K., Patra, N.K. & Singh, K. (2000) Performance of menthol mint (*Mentha arvensis*) genotypes to varying levels of nitrogen application under poplar based agro forestry system in Uttar Pradesh foot hills *Journal of Medicinal and Aromatic Plant Sciences*, 22 :447-449.
- Chauhan, H.S., Kamla Sing & Singh, H.B. (2000) Response of Palmarosa (*Cymbopogon martinii* var. Motia) to N and P fertilization in Uttar Pradesh foot hills of Himalayas. *Indian Perfumer*, 44: 61-64.
- Chinnamma, N.P. & Aiyer, R.S. (1988) Effect of fertilizers and harvests on palmarosa oil quality. *Indian Perfumer*, 32: 220-224.
- Chinnamma, N.P., Nair, E.V.G., Aiyer, R.S. & Saraswathy, P. (1988) Effect of fertilizers and harvest intervals on yield, nutrient composition and nutrient uptake of palmarosa. *Indian Perfumer*, 32: 278-287.
- Christopher, C., Rangaraju & KRISHNA, P.K. (1994) Importance of VAM. Intensive agriculture. May – June: 23-25.
- Dey, B.B. & Choudhuri, M.A. (1984) Effect of application of N, P and K on the growth and yield of essential oil and eugenol in *Ocimum sanctum* L. *Pafai Journal*, 6 : 20- 24.
- Earanna, N., Mallikarjuniah, R. R. Bagyaraj D. J. & Suresh, C. K. (2001) Response of *Coleus aromatics* to *Glomus fasciculatum* and other beneficial soil microflora. *Journal of spices and aromatic crops*. 10(2): 141-143.
- Farooqi, A.A., Devaiah, K.A., Dasharatha Rao, N.D., Vasundhara, M. & Raju, B. (1991) Effect of nutrients on growth, yield and essential oil content in Davana (*Artemisia pallens* Wall). *Indian Perfumer*, 35; 63-68.
- Farooqi, A.A., Devaiah, K.A., Vasundhara M. & Dasharatha Rao, N.D. (1990) Influence of planting season and plant density on growth, yield and essential oil in davana (*Artemisia pallens* Wall). *Indian Perfumer*, 34:274-277.
- Gowda, A.B., Krishnamurthy, K., Narayana, M.R., Rao, R.S.G., Chandrashekhar, G. & Puttanna, K. (1983b) Nutrient content and uptake studies in Java citronella (*Cymbopogon winterianus* Jowitt). *Mysore Journal of Agricultural Sciences*. 17 : 109-114.
- Gulati, B.C., Duhan, S.P.S. & Garg, S.N. (1978) Effect of nitrogen on the yield of herb, oil and quality of essential oil of *Ocimum basilicum*. *Indian Perfumer*, 22:53- 54.
- Gupta, M.L. & K.K. Janardhanan (1991) Mycorrhizal association of *Glomus aggregatum* with palmarosa enhances growth and biomass. *Plant and soil*.131(2): 261 – 264.
- Gupta, S., Arun Kumar & Khosla, M.K. (1992) Effect of growth regulators on biomass and oil yield of *Ocimum carnosum*. *Indian Perfumer*, 36: 27-32.
- Gupta, S.C. & Shahi, A.K. (1999) Spacing, nitrogen application and crop phenology studies on RRL-OC-11 (*Ocimum canum*). *Indian Perfumer*, 43: 95-101.
- Harshavardhan, P.G., Vasundhara, M., Srinivasappa, K.N., Biradar, S.L., Rao, G.G.E. & Gayithri, H.N. (2005) Effect of spacing and integrated nutrient management on biomass and oil yield in *Melissa officinalis*. *Indian Perfumer*, 49:349-354.
- Irulappan, I & V. Ponnusami (1982) Effect of nitrogen, phosphorus and potassium on growth and yield of patchouli. Proceedings of national seminar on Medicinal and Aromatic plants. Tamil Nadu Agricultural university. Pp. 141-144.
- Jadhav, S.G., Jadhav, B.B. & Apte, U.B. (2002) Influence of growth regulators on growth and oil content of patchouli. *Indian Perfumer*. 46: 287-289.
- Johri, A.K., Srivastava, L.J., Singh, J.M. & Rana, R.C. (1991) Effect of row spacings and nitrogen level on flower

and essential oil yield in German chamomile (*Matricaria chamomilla* L.) *Indian Perfumer*, 35:937-96.

Kaul, P.N., Rajeswar rao, B.R., Bhattacharya, A. K., Singh, K. & Singh, C.P. (1997) Effect of partial shade on essential oils of three Geranium cultivars. *Indian Perfumer*, 41: 1-4.

Kiruthika Devi (2002) Nutritional studies on root yield and quality of ashwagandha. M.Sc. (Hort.) thesis, HC & RI, TNAU.

Kothari, S.K., Ramesh, S. & Singh, K. (2003) Effect of harvesting frequency on oil yield and quality of lemon grass (*Cymbopogon flexuosus*) Cv. Krishna. *Indian Perfumer*, 47: 369-373.

Kothari, S.K., Singh, V. & Singh, K. (1987) Response of Japanese mint to varying levels of nitrogen application in Uttar Pradesh foot hills. *Indian Journal of Agricultural Sciences*, 57: 795-800.

Kothari, S.K., Singh, V.P. & Singh, U.B. (1996) The effect of row spacing and nitrogen fertilization on the growth, oil yield and composition of Japanese mint. *Journal of Medicinal and Aromatic Plant Sciences*, 18: 17-21.

Krishna Naik (1998) Effect of VA mycorrhizal and sloubilizing bacterium on growth and yield of java citronella at different sources of phosphorus. Ph.D Thesis. University of Agricultural Science, Bangalore.

Lokesh, D. (1979) Studies on nutrition and grafting of patchouli (*Pogostemon cablin*) on nematode resistant rootstock. M.Sc., (Agri) Thesis, University of Agricultural Sciences, Bangalore.

Madhaiyan, M., Santhanakrishnan, P. and Pragatheswari, D. (2000) Effect of orchid mycorrhizal fungi on the growth and nutrient status of *Vanilla planifolia*, Andr. *South Indian Hort.*, 49: 266-267.

Maheshwari, S.K., Gangrade, S.K. & Chouhan, G.S. (1991) Influence of planting geometry on irrigated palmarosa oil grass. *Indian Perfumer*, 35: 177-180.

Manjunatha, R., Farooqi, A.A., Vasandhara, M. & Srinivasappa, K.N. (2002) Effect of biofertilizers on growth, yield and essential oil content in patchouli. *Indian perfumer*. 46 (2): 97-104.

Munsi, P.S. (1992) Nitrogen and phosphorus nutrition response in Japanese mint cultivation. *Acta Horticulturae*, 306: 436-444.

Pal, S., Suresh Chandra, Balyan, S.S., Ajit Singh & Rao, B.L. (1992) Nitrogen requirement of new lemon grass strain -CKP 25. *Indian Perfumer*, 36: 75-80.

Panse, V.G. & Sukhatme (1957) Statistical methods for agricultural workers. ICAR publication, New Delhi. pp.125 – 148.

Rajadurai, K. R., K. Manivannan, M. Jawaharlal & A. Beaulah (2000) Effect of Azospirillum and VAM on growth characters of African marigold (*Tagetes erecta* L.) *South Indian Hort.*, 48(1-6): 83 -87.

Ramachandra, K. M., Vasundhara, M., Farooqi, A.A. & Srinivasappa, K.N. (2002) Evaluation of Patchouli (*Pogostemon cablin* Benth.) varieties in relation to different plant densities. *Indian Perfumer*, 46: 7-14.

Rao, B.R.R., Singh, S.P. & Rao, E.V.S.P. (1984) Effect of row spacings and nitrogen application on biomass yield, essential oil concentration and essential oil yield of bergamot mint (*Mentha citrata* Ehrh.) *Indian Perfumer*, 28:150-152.

Rao, P., E.V.S., Ganesha Rao, R.S., Narayana, M.R. & Ramesh, S. (1997) Influence of shade on yield and quality of patchouli (*Pogostemon cablin* Benth.)

Ratti, N. & Janardhanan, K.K. (1996) Response of dual inoculation with VAM and Azospirillum on the yield and oil content of palmarosa (*Cymbopogon martini* var. Motai). *Microbiological Res.*, 151(3): 325 -328.

Sadasakthi, A. (1986) Studies on the effect of planting system, nitrogen and phosphorus on growth and yield of marjoram (*Origanum majorana*). M.Sc. (Hort.). Thesis, Ac & RI, Madurai.

Saha, B.N., Baruah, B., Bardoli, D.N. & MATHUR, R.K. (1992) Prospects of growing patchouli (*Pogostemon patchouli* Pellet) in Arunachal Pradesh. Effect of nitrogen as its yield. *Indian perfumer*. 36(1) 57 -60.

Saha, B.N., Baruah, B.C. & Dutta, S.C. (1999) Response of nitrogen on herb and oil yield to BLI-1 strain of lemon grass (*Cymbopogon flexuosus*). *Indian Perfumer*, 43:156-158.

Samiullah, Varshney, A. K., Afridi, M.M.R.K, Mohammad, F. & AFAQ, S. H. (1988) Nitrogen requirement of lemon grass for optimum performance in western Uttar Pradesh. *Indian perfumer*, 32: 225-228.

Sastry, K.P., Karupaiyan, R., Mumtaz, S.H. & Sushilkumar (2000) Agronomic assessment of the oil yield in two chemotypes of geranium at different levels of nitrogen. *Journal of Medicinal and Aromatic Plant Sciences*, 22 :685-690.

Saxena, A. & Singh, J.N. (1996) Yield and nitrogen uptake of Japanese mint (*Mentha arvensis*) under various moisture regimes, mulch application and nitrogen fertilization. *Journal of Medicinal and Aromatic Plant Sciences*, 18: 477-480.

- Shalaby, A.S., Khattab, M.D., El-Gamassy, A. & EL-Gamassy, K. (1993) Cultivation of *Melissa officinalis* in Egypt; 1. Effects of fertilization, spacing and planting season. *Acta Horticulturae*, 331: 115-120.
- Sharma, S.N., Singh, A. & Tripathi, R.S. (1980b) Response of palmarosa to nitrogen, phosphorus, potassium and zinc. *Indian Journal of Agronomy*, 25: 719-723.
- Shelke, D.K. & Morey, D.K. (1978) Growth, yield and quality of Japanese mint (*Mentha arvensis*) as influenced by various levels of nitrogen and topping. *Journal of Maharashtra Agricultural University*, 3: 28-30.
- Shivalingappa, J. (1998) Influence of biofertilizers on growth, yield and essential oil content in tuberose (*polyanthus tuberosa*). M.Sc. Thesis, University of Agricultural Science, Bangalore.
- Singh, K., Singh, P.P., Beg, S.U., Kumar, D. & Patra, D.D. (2004) Effect of harvesting time and nitrogen application on essential oil yield and quality of a methyl chavicol rich cultivar of Indian basil (*Ocimum basilicum*). *Journal of Medicinal and Aromatic Plant Sciences*, 26: 293-296.
- Singh, M. & Dimri, B.P. (1997) Effect of N, P and K nutrition on herb and oil yield of *Artemisia annua* under semi arid tropical conditions. *Indian Perfumer*, 41: 45-46.
- Singh, M. & Ramesh, S. (2000) Effect of irrigation and nitrogen on herbage, oil yield and water use efficiency in rosemary grown under semi arid tropical conditions. *Journal of Medicinal and Aromatic Plant Sciences*, 22: 659-662.
- Singh, M. & Ramesh, S. (2002) Response of sweet basil (*Ocimum basilicum*) to organic and inorganic fertilizers in semi arid tropical conditions. *Journal of Medicinal and Aromatic Plant Sciences*, 24 : 947-950.
- Singh, M. & Ramesh, S. (2004) Effect of nitrogen and harvesting stage on oil yield of basil (*Ocimum basilicum* L.) in semi arid tropical climate. *Indian Perfumer*, 48:55-58
- Singh, M. & Sridhara, S. (2000) Direct and residual effect of intercrop rotation and nitrogen levels on performance of lemongrass. *Journal of Medicinal and Aromatic Plant Sciences*, 22: 260-263.
- Singh, M. (1996) Effect of irrigation and plant spacing on herb and oil yield of patchouli (*Pogostemon patchouli*) under semi-arid tropical conditions. *Journal of Medicinal and Aromatic Plant Sciences*, 18: 487-488.
- Singh, M. (1999) Effect of irrigation and nitrogen on herbage, oil yield and water use of lemon grass (*Cymbopogon flexuosus*) on alfisol. *Journal of Agricultural Sciences*, 132: 201-206.
- Singh, M. (2003) Effect of coconut and casuarina plants shade on growth, herbage and oil yield of aromatic crops under rainfed conditions. *Indian Perfumer*, 47:43-46.
- Singh, M., Baskaran, K., Kulkarni, R.N. & Ramesh, S. (2002) Comparative performance of lemon grass (*Cymbopogon flexuosus*) varieties at different levels of nitrogen. *Journal of Medicinal and Aromatic Plant Sciences*, 24 :50-52.
- Singh, M., Chandrashekhara, G. & Prakasa Rao, E.V.S. (1996) Effect of irrigation frequency and nitrogen on herb and oil yields of geranium, *Pelargonium graveolens* during summers season. *Journal of Medicinal and Aromatic Plant Sciences*, 18 : 495-495.
- Singh, M., Ramesh, S. & Sharma, S. (2002) Yield and oil quality of Rosemary (*Rosemarinus officinalis* L.) as influenced by nitrogen, irrigation, stages of harvesting in semi arid tropics. *Indian Perfumer*, 46: 15-20.
- Singh, R.L., Singh, N.P.J & gupta, K.C. (1981) Effect of levels of nitrogen application on the quality of essential oil of *Mentha* species. *Indian Perfumer*, 25: 78-83.
- Singh, U.B., Kothari, S.K. & Verma, B.S. (2002) Growth, yield and quality of mint species as intercrop in sugarcane under sub tropical climate of north Indian plains. *Journal of Medicinal and Aromatic Plant Sciences*, 24:60-64.
- Snedecor, G.W. & Cochran, W.G. (1967) Statistical methods, Oxford & IBH publishing Co. Calcutta.
- Sood, M., Singh, J.M., Rastogi, Srivastava, L.J. & Moresh Chand (1997) Effect of plant densities on German Chamomile (*Matricaria chamomilla* L.) *Indian Perfumer*, 41: 121-123.
- Sood, M., Singh, J.M., Srivastava, L.J. & Rana, R.C. (1996) Studies on the response of *Melissa officinalis* L. to nitrogen levels and row spacings. *Indian Perfumer*, 40:107-109.
- Sreeramulu, K.R. , Vishwanath shetty, Y. & Prabhakata Shetty, T.K. (1996) Effect of VA- Mycorrhizae on the growth of two important leafy vegetables. *Madras Agric. J.* , 83(6): 362-364.
- Subbaiah, B.V. & Asija, G.L. (1956) A rapid procedure for the estimation of available nitrogen in soils. *Current Science*, 25: 259-260.
- Umesha, K., Bojappa, K.M., Farooqi, A.A. & Suresh, N.S. (1990) Influence of plant density on growth and yield of clocimum (*Ocimum gratissimum* L.). *Indian perfumer*, 34: 173-175.
- Umesha, K., Bojappa, K.M., Suresh, N.S., Farooqi, A.A. & Jagannath, S. (1993) Herb, oil and eugenol yield of clocimum (*Ocimum gratissimum* L.) in relation to levels of nitrogen, phosphorus and potassium application. *Indian Perfumer*, 37: 311-314.



Vasundhara, M., Farooqi, A.A., Devaiah, K.A., & Shridharayya, M. (1992) Influence of some growth regulators on the growth, herbage and oil yield in marjoram (*Majorana hortensis Moench.*) *Indian Perfumer*, 36: 171-174.

Venugopal. C.K. (2006) Performance of Patchouli (*Pogostemon patchouli* Pellet) as influenced by nitrogen, spacing, shade, growth regulators and harvest techniques.

Ph.D. thesis. University of Agricultural Science, Bangalore

Vijayalalitha, S.J. & Rajasekaran LADA, R. (1996) Assimilation capacity of genotypes of patchouli (*Pogostemon patchouli* Pellet.) in open and shade. *Indian Perfumer*, 40: 113-114.