

## INTERNATIONAL JOURNAL OF ADVANCED BIOLOGICAL RESEARCH

© 2004-2014 Society For Science and Nature (SFSN). All Rights Reserved. www.scienceandnature.org

# LIPIDS AND FATTY ACIDS IN ACTIVE AND AESTIVATED HERMAPHRODITE PULMONATE GASTROPOD ACHATINA FULICA (BOWDICH)

\*Rakshit Abrol, S., <sup>a</sup>Bhattacharyya, D. K. & <sup>b</sup>Misra, K. K. \*Sanofi Oncology, 270 Albany Street, Cambridge MA, 02139, USA, <sup>a</sup>Department of Chemical Technology, Oil Technology Division, University of Calcutta, Kolkata 100 009, <sup>b</sup>Department of Zoology, R. B. C. College, Naihati 743 156, India. Corresponding author's email: misrakk47@gmail.com

#### ABSTRACT

Some pulmonate gastropods undergo aestivation for a long period. A hermaphrodite pulmonate gastropod, Achatina fulica, was selected for the study of some of the lipid components in active state and during aestivation of the snail. Thin layer chromatography and gas liquid chromatography were employed in the present study. Total lipid, phospholipid, cholesterol and triacylglycerol and the fatty acids of total lipid were studied in the digestive gland, ovotestis, mantle and foot of active and 7, 15, 30, 45, 60 and 90 days aestivated male and female A. fulica. The size of the albumen gland was the criteria of the identification of the sex and was confirmed through the present study. Percent lipid components in all the four organs exhibit clear male/female biasness. The digestive gland is the major lipid storage site over the other three organs and also plays important role in transportation of lipid to other organs during aestivation and maintains higher fatty acid percent than other organs. The other organs (albumen gland, mantle and foot) studied in this snail maintain a more or less steady lipid pool during aestivation. The fatty acid data also clearly indicate male/female dimorphism, and also emphasizes monoene to polyene conversion in this hermaphrodite gastropod.

KEYWORDS: Aestivation, Cholesterol, Fatty acid, Phospholipid, Snail, Triacylglycerol.

#### **INTRODUCTION**

Lipids are an integral part of molluskan tissues. Almost all data included in molluskan lipid study concern the entire organism and only scanty reports on the anatomical distribution of total lipids are available (Rakshit et al., 1997; Misra et al., 2002). Many mollusks store complex carbohydrates, in the form of glycogen, in the mantle and adductor muscle while lipids are accumulated in the digestive gland (Napolitano & Ackman, 1992). Contrarily several workers have proposed accumulation of lipids in other organs apart from digestive gland (Mitra & Sur, 1989; Wenne & Polak, 1989). Lipids from marine gastropods and bivalves are more extensively studied than freshwater or land counterparts as reviewed by a number of authors (Viarengo et al., 1994; Abad et al., 1995; Rakshit et al., 1997; Kraffe et al., 2004, 2005; Vale, 2010). Nevertheless, freshwater and land mollusks are important from the point of their evolutionary relationship within the sequence: marine mollusks-freshwater mollusks-land mollusks (Dembitsky et al., 1992). The basic types of lipids found to occur in mollusks include sterols, phospholipids, triacylglycerols, glycolipids to a lesser extent and fatty acids (Voogt, 1972). Major sterols occurring in gastropods include cholesterol,  $\Delta^{5,7}$  sterols, brassicasterol and  $\beta$  situaterol in different quantities (Bradshaw et al., 1991; Fried et al., 1998). Sterols in gastropods have been reported in two species of Cerethidea (Dutta et al., 1986; Misra et al., 1986). However it needs mention that almost all the data for sterol composition of mollusks concern the entire

organism (Frazer et al., 1997) and only few reports on the anatomical distribution of total lipids and sterols (Swift et al., 1980; Gordon & Collins, 1982; Klingensmith & Stillway, 1982; Stoilov et al., 1984) remain available. Dembitsky et al. (1992, 1993b,c, 1994) provide some information on phospholipid and fatty acid composition in pulmonates belonging to river Volga and Lake Baikal and also from some invertebrates (Dembitsky et al., 1993a). Triacylglycerols play an important role in reproductive strategy of mollusks and are incorporated into developing eggs (Barber & Blake, 1985; Napolitano et al., 1992) that are crucial for increasing larval survival and successful metamorphosis (Gallager & Mann, 1986). Dietary influence on sterol, triacylglycerols, and neutral lipids was indicated in studies with gastropod Biomphalaria glabrata fed with restricted diet (Conaway et al., 1996). Fatty acids have been studied in marine bivalves particularly because they serve as important source of polyunsaturated fatty acids (PUFAs) that are important from the standpoint of human nutrition and health (Johns et al., 1980; Jarzebski et al., 1986). Recently, studies on the lipids and fatty acids of fresh water mussels were also done (Ekin, 2012; Ekin et al., 2012). Fatty acid composition of freshwater gastropod mollusks of river Volga and Lake Baikal have been provided (Dembitsky et al., 1992, 1993 a,b,c). Biochemistry of PUFAs was studied in land snail Cepaea nemoralis by Van der Horst et al. (1973); Van der Horst & Oudejans (1976).and summarized in Ackman (1982). Zandee et al. (1980) indicated conversions of C<sub>18</sub> fatty acids to unsaturated  $\omega 6$  and  $\omega 3$  acids belonging to C<sub>20</sub> and

C<sub>22</sub> series in terrestrial herbivore snails. Aestivation is a simple physiological phenomenon exhibited by gastropods as a result of "low water content" which may be brought about either by "dryness of the environment" or "by the natural hydration cycle of the animal" (Swami & Reddy 1978). Chaki et al. (2008) studied activities of 13 metabolic enzymes in digestive gland, ovotestis and mantle of the active and aestivated Achatina fulica. The result showed that the enzymes activities were higher in the ovotestis during aestivation followed by digestive gland, while no remarkable change was observed in the mantle. The effects of aestivation on the phospholipidspecific fatty acid composition of mitochondrial membranes in the hepatopancreas of the terrestrial snail Cepaea nemoralis were investigated by Stuart et al. (1998). Some information about depletion of free fatty acids during aestivation in giant African snail Achatina achatina is provided by Umezurike & Iheanacho (1983). Pulmonates are hermaphrodites (Hyman, 1967; Parivar 1978), most species functioning as males in the early part of their life and females later on (Luchtel, 1972; Chétail & Fournié, 1986; South, 1992; Tomiyama, 1995). It is observed that Achatina fulica undergoes prolonged periods of aestivation and some of them lay eggs just after awakening at the onset of monsoon. It is obvious that both testis and ovary (within a single follicle) should not develop at a time because of hormonal regulation, and they mature at different age classes. This is evident through the development of albumen gland which is now considered as diagnostic marker of male-female distinction (Cunha et al., 1998). Mukai et al. (2004) confirmed that albumen gland is an accessory female reproductive structure of pulmonate (Egonmwan, 2007). This observation led the authors to think that prospective egg layers (females?) should exhibit a different physiological state than the others (males?). Thus a possible functional male/female distinction is possible on the basis of lipid and fatty acid storage. The present paper aims in obtaining a comprehensive knowledge on: a) lipid profiles in various organs viz., digestive gland, albumen gland, mantle and foot of active and aestivating hermaphrodite pulmonate gastropod, Achatina fulica, b) information about the lipid and fatty acid storage pattern, transportation and its utilization in these tissues during prolonged aestivation period based on male/female differentiation, and c) throw some light on the physiology of aestivation in A. fulica in relation to preparation for fecundity. The hypothesis of the present work is that male/female bias exists in A. fulica, which can be explained through the studies on lipid and fatty acid composition during active and aestivation.

## **MATERIALS & METHODS**

Collection of specimen: Healthy active snails, *Achatina fulica* (Bowdich), were collected from some selective fields around Calcutta. A total of 400 snails of an average size group (6.2 - 8.2 cm) were taken and 200 snails were allowed to aestivate. The snails selected for the experiment were more or less of similar in size and weight. The snails were induced to aestivate by placing them in an aestivation chamber under controlled temperature  $(20^{\circ}C \pm 2^{\circ}C)$  and humidity  $(45 \pm 10\%)$  coinciding the time of their natural

aestivation. The control specimens were kept active by providing them with leafy vegetables and water *ad libitum* throughout the experimental period. The control snails were also kept under controlled temperature  $(27^{\circ}C \pm 2^{\circ}C)$  and humidity  $(90 \pm 10\%)$  to minimize seasonal stress.

## Male/Female identification

Achatina fulica is a hermaphrodite snail and is very difficult to categorize them into male and female groups. In this study male and females were separated based on the size of their albumen glands during aestivation as it was observed that those with large albumen glands lay eggs (female) and those with smaller glands are males. Because it was observed that the snails having much larger albumen gland lay eggs after completion of aestivation and those with smaller albumen glands do not lay eggs. Thus during aestivation snails with larger albumen gland are supposed to act as females while others with small albumen gland remain as male (non-female).

## Collection of animal tissue

Control and aestivating gastropods were sacrificed at intervals of 7, 15, 30, 45, 60 and 90 days of experimentation. On each autopsy approximately 25 snails were sacrificed of which 10 males (with small albumen gland) and 10 females (with larger albumen gland) were considered. During autopsy, the snails were de-shelled and following tissues of the animal (*viz.*, digestive gland, albumen gland, foot and mantle) were carefully dissected out, excess mucous and water were wiped and weighed to a constant weight.

#### Extraction and analyses of lipid classes and fatty acids

Tissue lipid was extracted following the procedure of Bligh & Dyer (1959). The total lipid extracted from various tissues such as digestive gland, albumen gland, mantle and foot were weighed and residue was dissolved in known volume of fresh chloroform and nitrogen was bubbled through it (Kates, 1972). It was then stored in a freezer until further analysis.

Thin layer chromatoplates were prepared according to the general procedure described by Mangold (1969). Phospholipids were separated using solvent system of acetone-methanol-acetic acid-water (6: 8: 2: 1 v/v) (Rouser *et al.*, 1967). Neutral lipids were separated using hexane-diethyl ether-acetic acid (80: 20.1 v/v). All the lipids were identified by comparing Rf values with those of known standards and spraying with specific reagents.

Total cholesterol in tissue lipid was estimated according to Zlatkis et al. (1953). Phospholipids of tissue sample were determined by estimating phosphorus in the lipid according to standard procedure of Chen et al. (1956). Estimation of tissue triacylglycerol was carried out according to the standard procedure of Van Handel et al. (1957) after isolating pure triacylglycerol from total lipid. The fatty acid composition of the tissue lipids was determined by converting the lipids into methyl esters, separating the esters by thin layer chromatography and then analyzing it by gas liquid chromatography. Analysis of methyl ester was carried out by gas liquid chromatographic technique (Ghosh Choudhuri et al., 1983) in GC (Hewlett Packard Model 5890 A) with Flame ionization detector. The oven, injection port and the detector block temperatures used were 190°C, 230°C and  $240^{\circ}$ C respectively. The sample of methyl ester (0.1 - 1

ISSN 2250 - 3579

 $\mu$ g) was injected from a Hamilton syringe and the respective chromatogram was obtained. The fatty acids were identified by comparing the retention time of GLC grade standard saturated and unsaturated acid esters of varying chain length and unsaturation. The percentage composition of the component fatty acids was determined by measuring the areas of the peaks with the help of integrator (Hewlett Packard - 3390A).

#### RESULTS

The data on total lipid, phospholipid, cholesterol and triacylglycerol (TG) of active and aestivated *Achatina fulica* are presented in Table 1 (male) and Table 2

(female). Percent composition of total lipid in all the organs of active male of *A. fulica* are in following order: albumen gland (31.6%), digestive gland (31%), mantle (20.7%) and foot (16.7%) in comparison to female which shows; digestive gland (35.02), albumen gland (25.04), mantle (23.17) and foot (16.75) respectively (Figs. 1-4). Students 't' test indicate that the average total lipid of digestive and albumen glands in active male and female differ significantly at 0.5% level. In general mantle and foot contain lesser amount of total lipid in comparison to two other organs. Total lipid percent in digestive gland of male decreases at 45 and 90 days of aestivation; however, in female the variation is little.

**TABLE 1.** Total lipid, phospholipid, cholesterol and triacylglycerol (expressed in mg/g of wet tissues) in the different organs of male active and aestivating *Achatina fulica*. (mean±S.D.)

Days of aestivation				U	<b>v</b> ``	,		
Days of acstrvation		0 day	7 days	15 days	30 days	45 days	60 days	90 days
	Digestive gland	$39.08 \pm 8.93$	39.06±5.77	32.52±7.54	$25.65 \pm 3.41$	26.15±3.26	35.15±0.60	25.37±5.18
Total lipid	Albumen gland	39.86±1.03	36.17±3.89	25.27±8.70	29.79±6.82	29.65±2.81	31.67±4.01	36.73±3.09
	Mantle	26.11±1.94	20.06±3.45	21.63±2.62	$21.40\pm5.43$	19.23±2.28	23.67±3.11	20.27±4.04
	Foot	$21.07 \pm 4.61$	12.33±2.18	$10.88 \pm 1.49$	12.13±1.72	16.78±6.88	$14.07 \pm 3.43$	$15.44 \pm 0.87$
	Digestive gland	$14.27 \pm 2.82$	11.86±0.82	7.37±0.90	7.07±0.94	6.95±0.52	4.99±1.99	$2.82\pm0.91$
Phospholipid	Albumen gland	$14.05 \pm 1.54$	12.30±1.55	8.16±2.13	13.26±2.57	13.59±5.48	8.27±1.68	$9.95 \pm 2.40$
	Mantle	11.20±0.59	$10.77 \pm 1.88$	$11.38 \pm 6.57$	$11.43 \pm 1.64$	7.11±1.57	$5.22 \pm 3.08$	$4.86 \pm 1.68$
	Foot	6.27±0.23	5.85±1.59	5.24±0.53	$5.62 \pm 0.94$	4.94±0.55	3.02±1.69	3.13±1.57
	Digestive gland	$9.42 \pm 2.06$	$4.10 \pm 1.22$	$4.14 \pm 0.82$	4.23±0.38	4.41±1.29	$4.68 \pm 1.98$	$3.14 \pm 0.74$
Total cholesterol	Albumen gland	12.08±1.53	$5.45 \pm 2.05$	$3.45 \pm 1.89$	3.81±1.75	2.95±1.15	3.09±1.25	$2.65 \pm 0.83$
	Mantle	2.08±0.93	3.10±1.07	3.16±1.08	2.65±0.85	3.51±0.55	4.23±1.21	2.63±0.72
	Foot	2.67±0.92	$2.48\pm0.97$	2.53±1.31	$2.60 \pm 0.61$	$2.59 \pm 1.04$	$2.88 \pm 0.95$	2.15±0.99
	Digestive gland	3.61±0.95	$3.68 \pm 0.58$	2.43±0.32	3.13±0.18	$4.47 \pm 0.40$	2.99±0.56	2.23±0.10
Triacylglycerol	Albumen gland	$1.52\pm0.21$	$1.54\pm0.47$	$1.34\pm0.13$	2.19±0.09	$1.10\pm0.02$	$0.62\pm0.18$	$0.39 \pm 0.05$
	Mantle	$0.35 \pm 0.001$	$0.34 \pm 0.01$	$0.34 \pm 0.01$	$0.29 \pm 0.02$	$0.30 \pm 0.02$	0.34±0.09	$0.22 \pm 0.01$
	Foot	$0.24\pm0.01$	$0.24\pm0.11$	$0.09 \pm 0.00$	$0.32 \pm 0.01$	$0.06 \pm 0.02$	$0.14 \pm 0.01$	$0.20\pm0.06$

**TABLE 2.** Total lipid, phospholipid, cholesterol and triacylglycerol (expressed in mg/g of wet tissues) in the different organs of female active and aestivating *Achating fulica* (mean+S D)

	organs	of female acti	ve and aesti-	vating Achati	<i>ina fulica</i> . (m	ean±S.D.)		
Days of aestivation		→ 0 day	7 days	15 days	30 days	45 days	60 days	90 days
	Digestive gland	41.17±5.40	35.83±2.51	35.05±2.24	29.24±2.90	29.31±2.44	26.27±4.61	35.02±2.07
Total lipid	Albumen gland	17.51±2.26	26.47±3.50	21.09±2.79	$16.50 \pm 4.86$	$18.52 \pm 4.38$	$14.38 \pm 4.18$	$12.45 \pm 2.60$
	Mantle	26.99±1.49	20.37±2.24	22.20±5.04	19.94±4.66	20.13±6.14	20.98±3.50	21.11±3.94
	Foot	17.30±2.76	$15.03 \pm 3.75$	13.73±3.03	$12.03 \pm 1.84$	$15.72 \pm 4.85$	13.96±3.42	$14.12 \pm 1.25$
	Digestive gland	15.12±3.44	13.75±3.11	11.81±4.71	9.21±3.63	8.32±3.58	6.24±3.34	3.93±1.34
Phospholipid	Albumen gland	$6.05 \pm 2.16$	$8.62 \pm 1.88$	8.28±1.29	6.54±2.37	$5.20 \pm 1.78$	$5.86 \pm 1.40$	$4.44 \pm 1.82$
	Mantle	$13.83 \pm 2.40$	$9.48 \pm 3.18$	6.37±1.86	6.16±1.99	6.68±2.33	7.15±1.99	6.96±3.17
	Foot	6.43±2.59	5.76±1.41	4.83±1.08	4.50±1.05	$4.49 \pm 0.87$	3.62±1.32	$4.00 \pm 1.40$
	Digestive gland	5.90±1.17	4.21±1.74	$4.12\pm0.88$	4.97±0.54	3.99±1.39	4.78±1.61	3.11±0.78
Total cholesterol	Albumen gland	3.63±1.72	3.75±1.65	3.28±1.63	3.11±1.39	$3.74 \pm 0.85$	3.58±1.74	4.17±0.52
	Mantle	$3.03 \pm 1.58$	$2.50 \pm 1.07$	$2.15\pm0.94$	$2.24 \pm 1.10$	3.15±0.79	3.87±1.38	2.93±1.45
	Foot	2.75±1.37	$2.20\pm0.33$	$2.50\pm0.15$	2.31±0.92	2.41±1.45	$2.98 \pm 1.05$	2.53±1.14
	Digestive gland	$2.32 \pm 0.89$	$2.35 \pm 0.01$	$1.68\pm0.26$	$4.47 \pm 0.40$	3.17±0.50	$1.83 \pm 0.04$	$1.08\pm0.07$
Triacylglycerol	Albumen gland	$0.59 \pm 0.03$	$0.58 \pm 0.02$	$0.68 \pm 0.08$	$0.74\pm0.14$	$0.77 \pm 0.02$	$0.35 \pm 0.08$	$0.30\pm0.05$
	Mantle	$0.27 \pm 0.02$	$0.27 \pm 0.00$	$0.23 \pm 0.05$	0.31±0.01	$0.18\pm0.01$	0.19±0.23	$0.22 \pm 0.03$
	Foot	$0.09 \pm 0.00$	$0.11 \pm 0.00$	$0.08\pm0.00$	$0.24\pm0.02$	0.20±0.02	0.32±0.08	$0.17 \pm 0.11$
	80 -							
						glycerol		
	70 -			_	E Choles			
	60 -	<u>س</u>	-		La Phospi	lonpid		
	.g 50 - 🗱		হন ৫	> > >	2			
	. 40 - 🗱		n 🕅 🕯					
		887	* 99 8	3 2 13				
	30 - 8°			308		<		
	20 -				0 18 0			
	10 -					8 8		
	o _ 🖉							
	vfale	nale Male nale	Male nale	viale nale	nale Male male	nale		
		Fer Fer	Fer		Fer			
	0 d	lay 7 day	15 day Sex and da	30 day   45 d ays of aestivatio	ay   60 day on	90 day		

FIGURE 1. Percent composition of phospholipid, cholesterol and TG in total lipid of the digestive gland of active and aestivated *Achatina fulica* 

Lipids and fatty acids in active and aestivated hermaphrodite pulmonate gastropod



FIGURE 2. Percent composition of phospholipid, cholesterol and TG in total lipid of albumen gland of active and aestivated *Achatina fulica*.



FIGURE 3. Percent composition of phospholipid, cholesterol and TG in total lipid of mantle of active and aestivated *Achatina fulica.* 



FIGURE 4. Percent composition of phospholipid, cholesterol and TG in total lipid of foot of active and aestivated Achatina fulica

It is observed that the percentage of total lipid in albumen gland is always lower in female than that of the male during aestivation and the amount of phospholipid in the digestive gland of female is always higher than that of the male; though this amount decreases gradually with the progress of aestivation. The amount of phospholipid in albumen gland is higher in males than that of the female. The amount decreases little in aestivating snails of both the sex. Phospholipid content in both mantle and foot show slow decreases in amount as aestivation proceeds. Percent of phospholipid in total lipid also show similar pattern. Drop in phospholipid percent in 60 and 90 days aestivated male snail is lower than that of the female. The cholesterol level in digestive and albumen gland (it is obvious this is in the snail!) in both sexes reduces gradually throughout aestivation. Cholesterol content in mantle and foot shows fluctuation during aestivation. Percent of cholesterol in total lipid in all the organs is provided in Figs. 1-4. Table 3 tabulates ANOVA results of all the lipid classes during the progression of aestivation. Phospholipid/cholesterol ratio in digestive and albumen glands of control snail is always higher in female than that of the male (Table 4). However it is almost equal in the mantle and foot of control snails of both the sexes. During aestivation, this ratio is higher in female. Profile of TG in digestive gland of active and aestivating snail is more or less similar in both the sex and shows no significant difference. Percent of TG in total lipid fluctuates in digestive and albumen gland throughout the aestivation period. However, it remains fairly constant in mantle and foot (Figs. 3-4).

Classes of fatty acids and their composition in digestive gland, albumen gland, mantle and foot of active and aestivated snails of both sexes is represented in Tables 5-8 respectively. Pearson correlations coupled with p-value among fatty acid classes in digestive and albumen glands of active and aestivated snails show that male/female distinction is statistically significant (Table 9). Detailed account of all the fatty acid (percent composition w/w) classes in all the four organs with the ratios of saturated fatty acids (SFA), monoenes (M) and polyenes (P) is presented in Table 10.

**TABLE 3.** Analysis of variance of total lipid, phospholipid, cholesterol and triacylglycerol of all the organs of active and aestivated Achatina fulica of both sex

0 day						7 day					
	Sum of sq's	D.F.	Mean sq.	F(D.F./1)	Prob.		Sum of sq's	D.F.	Mean sq.	F(D.F./1)	Prob.
Regression	263.461	2	131.73	76.6371	0.08051	Regression	491.514	2	245.76	181.419	0.05243
Residual	1.7188	1	1.7188			Residual	1.3546	1	1.3546		
Total	265.179	3				Total	492.869	3			
15 day						30 day					
	Sum of sq's	D.F.	Mean sq.	F(D.F./1)	Prob.		Sum of sq's	D.F.	Mean sq.	F(D.F./1)	Prob.
Regression	243.657	2	121.829	697.482	0.02676	Regression	169.829	2	84.9144	49.3823	0.10012
Residual	0.17467	1	0.17467			Residual	1.71953	1	1.71953		
Total	243.832	3				Total	171.548	3			
45 day						60 day	·				
	Sum of sq's	D.F.	Mean sq.	F(D.F./1)	Prob.		Sum of sq's	D.F.	Mean sq.	F(D.F./1)	Prob.
Regression	106.253	2	53.1265	67.7434	0.08559	Regression	261.688	2	130.844	70.3964	0.08398
Residual	0.78423	1	0.78423			Residual	1.85867	1	1.85867		
Total	107.037	3				Total	263.547	3			
90 day											
	Sum of sq's	D.F.	Mean sq.	F(D.F./1)	Prob.						
Regression	250.242	2	125.121	2248.37	0.01491						
Residual	0.055649	1	0.05565								
Total	250.297	3									

TABLE 4. Phospholipid : cholesterol ratio in four organs of active and aestivating Achatina fulica of both sex.

Different organs		0 day	7 day	15 day	30 day	45 day	60 day	90 day
Digestive gland	Male	1.5:1	2.8:1	1.7:1	1.6:1	1.6:1	1.2:1	0.8:1
	Female	2.5:1	3.2:1	2.9:1	1.8:1	2.1:1	1.3:1	1.3:1
Albumen gland	Male	1.2:1	2.2:1	2.4:1	3.5:1	4.6:1	2.7:1	3.7:1
-	Female	1.7:1	2.4:1	2.5:1	2.1:1	1.4:1	1.6:1	1.1:1
Mantle	Male	5.4:1	3.5:1	3.6:1	4.3:1	2.0:1	1.2:1	1.8:1
	Female	4.6:1	3.8:1	3.0:1	2.8:1	2.2:1	1.8:1	2.4:1
Foot	Male	2.3 :1	2.3:1	2.1:1	2.1:1	1.9:1	1.1:1	1.4:1
	Female	2.3:1	2.6:1	1.9:1	1.9:1	1.9:1	1.2:1	1.6:1

22:4w3 22:5w6	20:5w3	20:3w3	18:4	18:3w3	Polyenes	22:1	18:1	16:1	15:1	14:1	24.0 Monoenes	18:0	18-0	16:0	14:0	Continented	T arry acrus	Fatty acide		24:2	22:5w6	22:4w3	20:5w3	20:4w6	20:3w3	18:4	18:3w3	Polyenes 18:2w6	22:1	18:1	16:1	15:1	14:1	24:0 Monoenes	0:81	17:0	16:0	15;0	14:0	Cotuestad	Fatty acids	
	$4.35\pm0.09$	8.07±0.15	'	$7.11\pm0.21$	5 N±71 A	$2.75 \pm 0.08$	$11.09 \pm 0.1$	$15.06 \pm 0.6$	$0.57{\pm}0.06$	$0.58{\pm}0.06$	,	1 C.0±C.7	0.36±0.06	17.98±0.1	$0.87 \pm 0.05$	INIDIC	Mala		TABLE 6.	I	$0.04{\pm}0.04$	1	$0.06{\pm}0.03$	$8.33{\pm}1.3$	$19.15 \pm 0.21$	$0.02 \pm 0.02$	$20.09\pm0.1$	$14.59 \pm 0.36$	$0.13{\pm}0.08$	$9.48{\pm}0.59$	$0.52{\pm}0.03$	$0.07{\pm}0.04$	$0.51{\pm}0.10$		11.48±0.67	$1.5\pm0.65$	$6.66 \pm 0.54$	I	$3.66 \pm 0.54$	Male	Control	TABI
$3.17{\pm}0.84$ $0.72{\pm}0.03$	$2.34\pm0.18$	9.99±0.07	$6.24 \pm 0.63$	1.94±0.72	0 6/+0 /5	$1.75 \pm 0.55$	7 11.11±0.20	2 13.74±0.24	$0.70{\pm}0.08$	$0.37 {\pm} 0.01$	0.72±0.31	0.02±0.21	0.32±0.08	8 19.40±0.10	$0.54\pm0.11$	LEIIIG	Empla	ontrol	Principal f	I	$0.03 \pm 0.01$	$0.31{\pm}0.03$	$0.01{\pm}0.0$	$9.54{\pm}0.20$	$18.32 \pm 0.51$	I	$21.02 \pm 0.74$	$10.63 \pm 0.57$	$0.04{\pm}0.03$	$13.94{\pm}1.27$	$0.84{\pm}0.07$	$0.73 \pm 0.08$	$0.61{\pm}0.1$	2.41±0.81	8.30±0.75	$4.61 \pm 0.95$	$6.59 \pm 0.31$	$0.70 \pm 0.32$	$3.16{\pm}0.65$	Female		LE 5. Princip
	$1.81\pm0.02$	6.66±0.06	ı	8.39±0.20	0 07±0 06	$2.45 \pm 0.29$	$0 13.66 \pm 0.64$	4 11.65±0.00	$0.49{\pm}0.2$	$1.52{\pm}0.34$	,	9.23±0.03		b 14.4±0.46	0.84±0.05	IVIALE	Mala	, T	atty acids c	I	$0.47{\pm}0.03$	I	$0.39{\pm}0.03$	$2.73 \pm 0.40$	$12.23 \pm 0.48$	$0.54{\pm}0.11$	$18.07 \pm 0.74$	$13.68 \pm 0.32$	$2.76 {\pm} 0.06$	$11.89{\pm}0.1$	$0.81{\pm}0.07$	$0.88 {\pm} 0.03$	$0.43{\pm}0.04$		7.86±0.15	$1.86 \pm 0.06$	$8.06 \pm 0.88$	I	$5.25 \pm 0.30$	Male	7 days	pal fatty acid
3.71±0.43	-	7.54±0.56	$1.98 \pm 0.03$	8.96±0.82	10 38±0 63	$1.80{\pm}0.09$	4 12.11±0.99	5 14.11±0.19	$1.39{\pm}0.17$	$1.54{\pm}0.66$	1.21±0.43	8.39±0.40		22.02±4.28	$0.65 \pm 0.03$	Temate	Emplo	fave +	of total lipid	I	$1.53 \pm 0.36$	1	$0.21{\pm}0.09$	$1.87{\pm}0.60$	$13.41 \pm 1.83$	$2.08 \pm 0.02$	$16.63 \pm 1.12$	$12.10\pm0.18$	$1.82 \pm 0.61$	$9.06{\pm}1.07$	$1.30{\pm}0.16$	$1.27 \pm 0.12$	$0.62{\pm}0.07$	1.12±0.92	6.61±0.35	3.57±0.59	$5.74\pm0.26$	I	$2.34{\pm}0.31$	Female		ls of total lip
	-	5.18±0.06	ı	6.35±0.30	0 06±0 05	ı	0 10.56±0.39	12.10±0.02	$0.14{\pm}0.02$	$0.13{\pm}0.1$	,	0.34±0.41		$14.26\pm0.08$	0.71±0.02	INICIE	Mala	15 J	of the albu	I	$0.88 \pm 0.03$		$0.38 \pm 0.09$	$1.36\pm0.45$	$12.38 \pm 0.42$	$0.36 \pm 0.07$	$10.27 \pm 0.98$	16.1±0.17	$2.54\pm0.37$	$16.05 \pm 0.33$	$4.33 \pm 0.46$	$1.63 \pm 0.41$	$0.48 \pm 0.04$		$8.81\pm0.16$	$3.31\pm1.04$	$14.27 \pm 0.62$	I	$3.69 \pm 0.22$	Male	15 days	id of the dig
1 1	-	4.80±0.09	'	8.18±0.56	10 38±0 03	$2.04{\pm}0.16$	$13.63{\pm}0.05$	$13.06{\pm}0.18$	$0.56 \pm 0.02$	$0.53{\pm}0.04$	0.2010.32	0 00±0 20		$20.40\pm1.30$	$0.91 \pm 0.02$	Tennale	aya Eamala	ave	ımen gland			1	-	1.51±0.61	$16.34{\pm}1.02$	5.02±0.53	16.96±0.50	14.45±0.93	$0.75 \pm 0.14$	$15.04 \pm 1.53$		0.76±0.14 (	$0.48 \pm 0.07$	0.98±0.2 -	5.4±0.28	3.51±0.30	8.57±0.41		4.49±0.10	Female		estive gland
1.35±0.04 -	-	7.92±0.04	ı	6.12±0.08	2 N0±N 17	I	$10.19 \pm 0.24$	$9.64{\pm}0.01$	$0.67{\pm}0.04$	$0.62{\pm}0.01$	,	9.000.09		19.36±0.02	$0.99 \pm 0.02$	INTOTE	Mala	sp 05	of active a	-		1	).27±0.2 -	$2.47 \pm 0.27$ 1	15.78±2.36 1	1.29±0.69 2	$3.43 \pm 0.98$ 1	15.0±1.63 1	l.94±0.81 -	14.21±1.59 1	2.99±1.29 -	).84±0.15 1	l.16±0.53 1	- -	10.52±1.36	2.12±0.56 2	12.61±2.91 1		7.05±1.92 4	Male F	30 days	of active an
	-	12.99±0.45	ı	7.45±0.25	11 01±0 21	$0.98 \pm 0.25$	$12.46 \pm 0.32$	$6.26 \pm 1.63$	$1.11 \pm 0.21$	$0.34{\pm}0.01$	1.12±0.04	0.41±0.23		17.99±1.85	0.93±0.25	TEIII	ya Earrala	Ve	nd aestivat	2		1		.31±0.17 1	$1.55 \pm 1.82$ 1	.78±0.67 4	5.85±0.58 9	5.80±2.52 1	3	7.51±1.00 1	4	.08±0.27 0	.02±0.22 1	).54±0.12 -	.45±0.33 /	1.98±2.26 2	0.73±2.92 8	1	.44±3.61 3	emale N		d aestivated
1.09±0.03 -		4.70±0.09	'	7.08±0.06	2 18±0 08	ı	$11.76 \pm 0.14$	$12.14 \pm 0.15$	$1.91{\pm}0.03$	$1.78{\pm}0.11$	,	1.19±0.17		$16.62\pm0.1$	$3.09 \pm 0.17$	INIG	Mala	45 d	ed male an	.91±0.89 -		.71±0.91 -		.73±0.51 1.	4.49±0.51 1	.23±1.03 4	.16±0.06 1.	6.50±0.44 1	.08±0.14 -	5.52±0.37 1	.96±0.16 -	.96±0.08 0.	.13±0.18 0.	0	.10±0.67 4.	.94±0.80 1.	.22±0.62 9	ı	.35±0.45 5	fale F	45 days	male and fe
1.18±0.16 -	- -	7.63±0.03	ı	5.44±0.41	11 12±0 12	$0.32 \pm 0.01$	$9.67 \pm 0.31$	$1.93{\pm}0.05$	$2.14\pm0.24$	$0.81 \pm 0.02$	2.30±0.03	2.20±0.00		$12.71\pm0.17$	$1.28 \pm 0.06$	TEITIAE	Tamala	ave	d female A	I		1		.61±0.10 5	$3.34 \pm 0.51$ 1	$.25\pm0.12$ 4	$4.55 \pm 0.11$ 1	6.65±0.36 1	1	$4.43 \pm 1.10$ 1	0	.71±0.11 0	.41±0.11 0	- cc.0±8/.	.55±0.15 /	$.69\pm0.44$ 1	$.62\pm0.25$ 1	I	.78±0.14 1	emale N		male Achat.
1.05±0.06 -	-	8.70±0.03	ı	-	10.05±0.13	I	$13.57 \pm 0.51$	17.78±0.24	$1.23{\pm}0.05$	$1.60{\pm}0.08$	,	8.3/±0.27	-	15.04±3.41	$1.16\pm0.1$	INTAIC	Mala	60 /	chatina fuli	-	1	1		.07±1.62 3	$2.93 \pm 4.99 $ 1	.23±1.17 5	1.62±3.46 1	7.83±0.70 1	.44±0.01 -	5.02±2.01 1	.73±0.11 -	.80±0.09 0	.49±0.34 0	2	.88±0.54 9	$.15\pm0.14$ 1	$3.64\pm0.11$ 9	1	.57±0.05 2	Iale F	60 days	ina fulica. (
1.88±0.06 -		8.52±0.10	ı	5.24±2.93	17 6+08 0	$1.04{\pm}0.15$	$11.95\pm1.28$		$0.55 \pm 0.22$	$0.35 \pm 0.13$	1.70±0.00	4.81±1.20	-	17.92±1.49	$1.77\pm0.72$	LEIIIT	Ermolo	AVG	ca. (mean±	2	.10±0.02 -	_		.49±1.20 -	4.77±0.54 1	.38±1.89 7	6.89±0.68 8	7.21±1.03 1	2	3.32±1.16 1	1	.41±0.22 0	.48±0.04 0	- 1C.0±20.	.36±2.84	.34±0.38 1	.41±2.51 5	1	.71±0.08 4	emale N		mean±S.D.)
1.14±0.15 -		5.89±0.89	ı	-	6 60±0 00	ı	8 13.25±2.0	$16.0\pm2.24$	$1.03\pm0.14$	$5.47{\pm}1.11$	,	/.20±0.42		9 15.38±1.8	1.53±0.49	IVIALE	Mala	90 dave	S.D.)	$2.49 \pm 1.48$		.49±0.83			$7.23\pm2.34$	.48±0.86	2.63±3.03	$3.52 \pm 0.77$	$2.49 \pm 0.80$	$3.81{\pm}1.00$	$.32{\pm}0.03$	).63±0.17	1.49±0.03		8C.0±C/.7	$1.99 \pm 0.81$	$5.88 \pm 0.29$		.52±2.33	Viale	90 days	
1.60±0.51 -	-	8.30±0.51	ı	$7.26 \pm 1.77$	17 2+22 17	ı	$3 15.02 \pm 1.28$	ı	$0.38{\pm}0.03$	$0.38{\pm}0.19$	2.37王0.42	4.33±2.02		9 16.96±2.12	$0.90 \pm 0.08$	TEIIIAIC	Empla		29	92	$1.28{\pm}0.41$			$5.16\pm0.66$	$9.27{\pm}1.1$	$4.77 \pm 0.49$	$12.84 \pm 0.73$	$13.78 \pm 1.43$		$14.10{\pm}1.0$		$0.28 \pm 0.06$	$0.14{\pm}0.04$	5.96±1.56	5.99±1.60	$1.23\pm0.24$	$11.06\pm 2.73$	1	$3.56 \pm 1.80$	Female		

0MC:77	22:4w3	20:5w3	20:4w6	20:3w3	18:4	18:3w3	18:2w6	Polyenes	22:0	18:1	16:1	15:1	14:1	INTOLIOEUES	24:0	24.0	18:0	17:0	16:0	14:0	Saturated		Fatty acids		22:5w6	22:4w3	20:5w3	20:4w6	20:3w3	18:4	18:3w3	18:2w6	Polyenes	22:0	18:1	16:1	15:1	14:1	Monoenes	24:0	18:0	17:0	16:0	14:0	Saturated		Fatty acids
$0.20\pm0.08$	$7.19 \pm 0.57$	$0.64{\pm}0.15$	$2.30 \pm 0.45$	$12.16 \pm 0.70$	$7.30 \pm 0.35$	$4.71 \pm 0.60$	$9.95 \pm 0.24$		$3.06 \pm 0.84$	$11.63 \pm 0.58$	$1.52\pm0.34$	$1.99\pm0.07$	0.56±0.28	0 2 . 0 00	/.U±U.Uo	70.000	8.44+0.42	$4.12 \pm 0.30$	$8.54{\pm}0.56$	$0.31{\pm}0.09$		Male	Control		$0.21 \pm 0.02$	$6.26 \pm 1.61$	$1.29{\pm}0.13$	$1.92 \pm 0.37$	14.84±0.33	$6.51\pm2.10$	$3.92{\pm}1.45$	$9.03 \pm 0.79$		$5.75 \pm 0.93$	$8.60 \pm 1.22$	$0.47{\pm}0.06$	$0.71 \pm 0.26$	$0.34{\pm}0.10$		$6.06 \pm 1.62$	$6.54{\pm}0.61$	$0.90 \pm 0.20$	$18.08 \pm 0.81$	$1.01{\pm}0.85$		Male	Contro
2.08±0.97	$4.11 \pm 0.18$	$5.52 \pm 0.54$	$1.80 \pm 0.96$	$8.78{\pm}1.10$	$5.17 \pm 0.25$	$3.03 \pm 0.86$	$11.65{\pm}1.09$		$3.43{\pm}1.11$	$17.97 \pm 0.07$	$1.41{\pm}0.69$	$1.7/\pm0.09$	$0.33\pm0.14$	000011	1.20世0.23	1 20-0 25	10.50+0.61	$1.43 \pm 0.57$	$6.93 \pm 0.68$	$0.33 \pm 0.14$		Female		TAB]	$0.45 \pm 0.08$	$5.36 \pm 1.30$	$1.52 \pm 0.67$	$2.85 \pm 0.91$	3 15.30±0.3€	$9.0{\pm}1.53$	$5.64{\pm}1.24$	$11.50{\pm}1.64$		$7.17 \pm 1.17$	$11.79 \pm 1.09$	$0.35 \pm 0.24$	$0.83 \pm 0.14$	$0.41\pm0.17$		$4.11 \pm 0.99$	8.08±0.73	$0.50 \pm 0.08$	1 8.01±1.65	$0.24{\pm}0.07$		Female	51
	$4.02 \pm 0.10$	$0.57 \pm 0.14$	$1.74{\pm}0.48$	$10.16 \pm 0.56$	$9.23 \pm 0.13$	$6.79 \pm 1.23$	$9.54{\pm}0.23$		$5.41 \pm 0.68$	$16.40 \pm 0.60$	$1.34{\pm}0.38$	0.86±0.05	0.65±0.19		4.98±0.40	4 00 0 46	8.0+0.70	$1.78{\pm}0.98$	$8.42 \pm 0.44$	$1.11{\pm}0.18$		Male	7 days	LE 8. Princi	$0.29 \pm 0.08$	, <b>'</b>	ı	$2.91 \pm 0.25$	$11.49 \pm 1.60$	$8.60 \pm 0.61$	8.77±0.93	· 10.79±0.59		$6.92 \pm 0.37$	11.43±0.68	$0.61{\pm}0.20$	$1.45 \pm 0.33$	$1.30 \pm 0.07$		$2.38 \pm 0.50$	$7.90 \pm 0.26$	$1.95 \pm 0.40$	$11.95 \pm 0.12$	$3.84{\pm}0.56$		Male	7 days
	$2.34{\pm}0.32$	$3.79 \pm 0.59$	$1.06 \pm 0.06$	$7.63 \pm 0.56$	$7.27\pm0.79$	6.85±0.75	$8.94{\pm}0.68$		$3.02 \pm 0.54$	$13.37 \pm 0.59$	$0.66 \pm 0.55$	2.40±0.53	0.39±0.28	00000	1.23±0.24	105-004	8 30+0.60	$2.50\pm0.57$	$6.70 \pm 0.66$	$1.19{\pm}0.28$		Female		pal fatty acid	$0.28 \pm 0.11$	$3.43 \pm 0.54$	$1.12{\pm}0.01$	$1.86 \pm 0.59$	$13.22 \pm 0.38$	8.75±0.64	$7.35 \pm 0.50$	$11.06 \pm 0.54$		$7.13 \pm 0.23$	$9.92{\pm}0.69$	$0.82 \pm 0.45$	$4.14{\pm}0.56$	$0.79 \pm 0.10$		$4.29{\pm}0.56$	$6.94{\pm}0.13$	$2.01 \pm 0.51$	$7.78 \pm 0.60$	$1.70{\pm}0.60$		Female	
	$7.51{\pm}1.09$	$1.15 \pm 0.06$	$1.68 \pm 0.76$	$9.94{\pm}0.06$	$7.34{\pm}0.58$	$8.93 {\pm} 0.95$	$8.39 \pm 0.43$		$3.97{\pm}0.10$	$15.29 \pm 0.45$	$5.27 \pm 0.40$	$0.54\pm0.23$	1.95±0.73	100.00	4.9/±0.0/	4 07 0 67	4.43+0.62	$1.02{\pm}0.01$	$13.15 \pm 0.57$	$0.96 \pm 0.55$		Male	15 days	ds of total lip		'	·	$1.43 \pm 0.11$	$15.02 \pm 0.11$	$8.84 \pm 0.65$	$9.06 \pm 0.18$	$14.06 \pm 0.13$		$7.76 \pm 0.31$	$14.55 \pm 0.44$	$0.29{\pm}0.09$	$0.76 \pm 0.06$	$0.40 \pm 0.11$		$3.19{\pm}0.28$	$8.82 \pm 0.71$	$2.36 \pm 0.06$	$10.08 \pm 0.52$	$1.52{\pm}0.11$		Male	15 days
	$2.42\pm0.13$	$3.34{\pm}0.56$	$1.15 \pm 0.14$	$8.14{\pm}0.52$	$10.10 \pm 0.60$	$6.68 \pm 0.23$	$9.43 \pm 0.52$		$4.43 \pm 0.54$	$19.21 \pm 0.53$	$1.19{\pm}0.25$	$3.31\pm0.58$	$0.42\pm0.11$	0 10 0 11	2.32世0.02	0.00101000	8.70+0.60	$6.09{\pm}0.69$	$10.02 \pm 0.54$	$1.09{\pm}0.55$		Female		pid of the fo		$2.89 \pm 0.54$	I	$1.25{\pm}0.58$	$8.08 {\pm} 1.20$	$3.08 \pm 0.61$	$8.12 \pm 0.54$	$10.83 {\pm} 0.59$		$7.27{\pm}0.39$	$14.31\pm2.11$	$1.88 {\pm} 0.75$	$1.39{\pm}0.08$	$1.31 \pm 0.39$		$8.43 \pm 1.02$	$11.44 \pm 0.54$	$1.50 \pm 0.45$	$11.38{\pm}1.69$	$1.94{\pm}0.67$		Female	
	$2.75 \pm 0.20$	$0.36 {\pm} 0.03$	$1.07{\pm}0.06$	$9.19{\pm}0.11$	$8.74{\pm}0.33$	$6.17{\pm}0.58$	$7.75{\pm}1.72$		$3.44{\pm}0.24$	$15.06 \pm 1.76$	$2.25 \pm 0.98$	2.3/±0.85	2.60±0.60		4./1±0./9	4 71 0 70	11.75+2.31	$2.13 \pm 0.32$	$11.90\pm0.43$	$1.34{\pm}0.32$		Male	30 days	ot of active a			ı	$1.07{\pm}0.77$	$16.21\pm0.24$	$9.48{\pm}0.64$	$7.51\pm0.18$	$9.94{\pm}1.16$		$5.27{\pm}0.03$	$11.87 \pm 0.56$	$1.32{\pm}0.24$	$1.32{\pm}0.24$	$0.30 \pm 0.10$		$1.50 \pm 0.70$	$11.89 \pm 0.82$	$4.00{\pm}2.17$	$8.37{\pm}1.85$	$1.00{\pm}0.35$		Male	30 days
	$3.17 \pm 0.56$	$3.98 \pm 0.78$	$1.70{\pm}1.05$	$7.85 \pm 0.43$	$10.21 \pm 0.28$	$6.90{\pm}0.84$	$9.33 \pm 0.05$		$4.09 \pm 0.60$	$16.73 \pm 1.84$	$2.40\pm0.59$	$3.0' \pm 1.18$	0.14±0.05	011.005	$1.07\pm0.14$	1 07 0 14	9.69+0.58	$6.90 \pm 0.27$	$11.45 \pm 3.34$	$1.24{\pm}0.20$		Female		und aestivate		$1.89{\pm}0.43$	ı	$2.40 \pm 0.29$	$15.95{\pm}1.09$	$9.98 \pm 0.66$	$5.04\pm2.36$	$10.57 \pm 1.96$		$6.39 \pm 2.50$	$11.21\pm0.77$	$2.14{\pm}1.20$	$2.20{\pm}0.26$	$0.30 \pm 0.14$		$6.87 \pm 1.70$	$9.40 \pm 0.43$	$3.50{\pm}1.52$	$10.74 \pm 0.48$	$1.84{\pm}0.34$		Female	
	$2.16 \pm 0.22$	ı	$1.32{\pm}0.10$	$9.86 {\pm} 0.56$	$7.62 \pm 0.72$	$5.64{\pm}0.71$	$10.89 \pm 0.77$		$3.95 \pm 0.66$	$18.30 \pm 0.44$	$1.60 \pm 0.44$	1./0±0.05	1.70±0.01	0 20.001	J.27±0.33	C 07 0 00	8 44+0 65	$1.76{\pm}0.49$	$12.58 \pm 0.58$	$2.33 \pm 0.40$		Male	45 days	ed male and		$2.32 \pm 0.38$	ı	$1.98{\pm}0.06$	$17.40\pm0.45$	$9.82 \pm 0.09$	$6.41 \pm 0.52$	$10.60 \pm 0.35$		$6.76 \pm 0.91$	$10.44 \pm 0.54$	$0.53{\pm}0.05$	$2.05 \pm 0.16$	$0.43 {\pm} 0.05$		$6.19{\pm}0.22$	$6.35 \pm 0.40$	$3.46 \pm 0.56$	$7.66 \pm 0.48$	$3.26 \pm 0.33$		Male	45 days
	$1.56 \pm 0.31$	$2.91 \pm 0.55$	$1.21 \pm 0.02$	$12.05 \pm 0.73$	$9.05 \pm 0.77$	$5.41 \pm 0.61$	$10.73 \pm 0.60$		$3.99 \pm 0.07$	$13.39 \pm 0.60$	$1.24{\pm}0.07$	3.51±0.54	0.50±0.08	0 50.000	2.30±0.04	0.05-0.04	6.59+0.97	$3.73 \pm 0.44$	$9.42{\pm}0.55$	$2.90{\pm}0.54$		Female		female Achu		$1.74{\pm}0.09$	$2.51 \pm 0.54$	$4.08 \pm 0.16$	$16.54 \pm 0.55$	$10.47 \pm 0.49$	$5.28 \pm 0.35$	$10.31 \pm 0.36$		$4.08{\pm}0.16$	$10.59 \pm 0.52$	·	$4.70 \pm 0.54$	$0.98{\pm}0.06$		$6.63 {\pm} 0.62$	$5.96 \pm 0.72$	$3.86 {\pm} 0.71$	$10.80 \pm 0.08$	$1.09{\pm}0.54$		Female	
	$2.79 \pm 0.32$		$1.83 {\pm} 0.51$	$11.86 \pm 3.96$	$6.86{\pm}2.06$	$6.55 \pm 0.87$	$10.14{\pm}1.52$		$4.91 \pm 0.36$	$18.26 \pm 1.48$	$1.31 \pm 0.33$	$0.94 \pm 0.04$	0.55±0.52	0	/.14±0.14	714.014	9.12+0.88	$2.26 \pm 0.21$	$12.11 \pm 1.77$	$1.00{\pm}0.52$		Male	60 days	utina fulica.		$1.85 \pm 0.65$	ı	$1.58{\pm}0.71$	$17.46\pm0.71$	$9.10 \pm 0.35$	$7.24\pm0.59$	$10.64{\pm}0.08$		$6.77 \pm 1.79$	$11.91 \pm 0.89$	$0.39{\pm}0.15$	$1.28{\pm}0.07$	$0.75 \pm 0.45$		$6.35 \pm 0.09$	$7.58 \pm 3.66$	$2.0\pm0.84$	$7.66 \pm 0.14$	$1.92{\pm}0.28$		Male	60 days
,	$1.60 \pm 0.41$		ı	$9.13{\pm}1.58$	$8.86 \pm 0.05$	$6.03 \pm 0.56$	$11.12\pm0.18$		$2.14{\pm}0.06$	$18.07 \pm 3.78$	$1.63 \pm 0.05$	$0.82 \pm 0.33$	0.49±0.23	0 10 .0 22	0.U/±0.00	E 07 0 E2	10.43+1.23	$1.69{\pm}0.34$	$9.31 \pm 3.57$	$1.76 \pm 0.62$		Female				$1.79{\pm}0.09$	·	$1.74{\pm}1.39$	$14.51\pm1.39$	$9.67 \pm 2.10$	$5.78 \pm 0.89$	$8.71 \pm 3.05$		$5.76 \pm 1.53$	$14.55 \pm 0.70$	'	$1.79{\pm}0.08$	$0.76{\pm}0.14$		$7.06{\pm}1.09$	$7.82 \pm 1.93$	$2.32 \pm 0.88$	$10.68 \pm 0.16$	$3.13 \pm 0.31$		Female	
	$2.68 \pm 0.23$	ŗ	$1.36{\pm}0.05$	$12.57 \pm 2.13$	$9.92 \pm 1.12$	$5.70 \pm 1.14$	$11.67{\pm}0.84$		$3.73 \pm 1.32$	$17.04 \pm 0.73$	$0.56 \pm 0.14$	$0.50\pm0.09$	$0.48\pm0.44$	0 10 0 11	0.92±0.00	600-050	8 56+1.08	$0.97{\pm}0.64$	$7.46 \pm 0.05$	$1.50 \pm 0.42$		Male	90 days			$1.95 \pm 0.19$	ı	ı	$18.86 \pm 0.06$	$9.07 \pm 0.38$	$5.73 \pm 0.38$	$12.15 \pm 1.43$		$7.42{\pm}0.10$	$14.31 \pm 2.06$	$2.12 \pm 0.42$	$0.27{\pm}0.05$	$0.29{\pm}0.02$		$6.52{\pm}0.86$	$7.40{\pm}2.23$	$0.61 \pm 0.35$	$8.29 {\pm} 1.99$	$1.0\pm0.30$		Male	90 days
,	$3.04 \pm 0.27$		ı	$13.94{\pm}1.07$	$11.80\pm0.12$	$5.76 \pm 1.16$	$10.70 \pm 1.13$		ı	$18.68 \pm 0.82$	$0.67 \pm 0.31$	$1.11\pm0.18$	$0.22\pm0.02$		0.00±0.40	C.CC_2.10	8.83+2.40	$1.70\pm0.61$	$9.19 \pm 1.12$	$1.21 \pm 0.32$		Female				$2.15 \pm 0.16$	ı	·	$13.99 \pm 0.32$	$11.13 \pm 3.46$	$7.93\pm2.97$	$11.38 \pm 0.92$		$10.73 \pm 2.50$	$14.85 \pm 2.08$	'	$0.79 \pm 2.50$	$0.24{\pm}0.13$		$5.63 \pm 0.37$	$6.66 \pm 1.79$	$0.81 \pm 0.11$	$824 \pm 0.90$	$2.01 \pm 0.58$		Female	

TABLE 7. Principal fatty acids of the total lipid of mantle of active and aestivated male and female Achatina fulica.

	DgMs	DgFs	AlMs	AlFs	DgMm	DgFm	AlMm	AlFm	DgMp	DgFp	AlMp
DgFs	0.076										
	0.871										
AlMs	0.245	0.439									
	0.596	0.324									
AlFs	-0.468	-0.140	-0.313								
	0.289	0.765	0.494								
DgMm	0.373	0.056	0.481	0.395							
	0.409	0.906	0.275	0.381							
DgFm	0.508	-0.483	-0.391	0.027	0.331						
	0.244	0.272	0.385	0.953	0.469						
AlMm	0.094	0.431	-0.145	0.175	-0.205	-0.412					
	0.841	0.335	0.756	0.708	0.660	0.358					
AlFm	-0.019	-0.631	-0.612	0.054	-0.098	0.801	-0.566				
	0.967	0.128	0.144	0.908	0.834	0.03	0.185				
DgMp	-0.128	-0.358	-0.350	-0.423	-0.824	-0.29	0.246	-0.036			
	0.784	0.430	0.441	0.344	0.023	0.528	0.595	0.939			
DgFp	-0.288	-0.826	-0.476	0.072	-0.405	0.025	-0.129	0.258	0.717		
	0.531	0.022	0.280	0.879	0.367	0.958	0.782	0.576	0.070		
AlMp	-0.168	-0.584	-0.222	-0.064	0.031	0.557	-0.932	0.780	-0.115	0.300	
	0.718	0.169	0.633	0.891	0.948	0.194	0.002	0.039	0.805	0.513	
AlFp	0.145	0.414	0.733	-0.526	-0.070	-0.698	0.172	-0.823	0.270	-0.076	-0.434
	0.757	0.355	0.061	0.225	0.881	0.081	0.712	0.023	0.557	0.871	0.331

<b>TABLE 9.</b> Correlations among three fatty	y acid classes in digestive and	1 albumen glands of	active and aestivating Achatina
	<i>fulica</i> of both sex.		

Cell contents : Pearson correlation P- value Dg = digestive gland Al = albumen gland M = maleF = female s = saturated fatty acid m = monoene fatty acid p = polyenefatty acid

**TABLE 10.** Principal fatty acid of total lipid composition (% w/w) in all the organs of active and aestivating Achatina fulica of both sex

Days of		Dige	stive gland		Alb	umen gland			Mantle			Foot			Total		
acstivation	Sex	Saturated	Monoene	Polyene	Saturated	Monoene	Polyene	Saturated	Monoene	Polyene	Saturated	Monoene	Polyene	Saturated	Monoene	Polyene	SFA:M:P ratio
0 day	Male	24.20	11.12	64.68	30.05	33.87	36.07	35.26	17.17	47.48	30.40	20.29	48.96	119,91	82.45	197.21	1.45:1:2.39
NR3178	Female	23.56	16.30	60.14	29.77	28.42	41.81	22.53	22.11	55.35	22.93	28.02	49.04	100.59	94.85	206.34	1.06:1:2.17
7 day	Male	26.20	19.08	54.73	28.05	34.40	37.93	30.27	23.45	46.28	26.70	27.08	46.22	111.22	104.01	185.16	1.07:1:1.78
2020-022	Female	23.84	21.00	58.85	32.38	31.05	35.56	24.54	24.62	50.84	25.67	24.25	48.77	106.43	100.92	190.02	1.05:1:1.88
15 day	Male	31.06	25.85	43.09	30.21	32.21	37.58	26.46	24.21	49.33	25.53	28.10	46.36	113.26	110.37	176.36	1.03:1:1.60
1	Female	23.71	2.23	56.05	37.26	32.11	30.63	36.47	27.50	36.00	28.71	29.13	42.08	126.15	108.99	164.76	1.16:1:1.55
30 day	Male	33.41	21.87	44.72	33.16	28.85	37.99	30.25	22.70	47.04	34.00	24.48	38.49	130.82	97.90	166.24	1.34:1:1.70
	Female	29.83	16.21	53.96	32.10	23.86	44.04	31.50	22.36	46.09	30.37	26.45	43.17	123.80	88.88	187.26	1.39:1:2.11
45 day	Male	22.05	26.18	51.77	34.47	34.58	30.96	28.67	20.97	50.36	32.35	24.74	39.92	117.54	106.47	173.01	1.10:1:1.62
	Female	25.37	17.61	57.03	30.59	21.01	48.40	27.92	20.58	51.50	27.60	24.99	47.40	111.48	84.19	204.33	1.32:1:2.43
60 day	Male	24.76	19.82	55.42	30.28	41.79	27.92	27.00	22.33	50.67	32.40	26.60	41.00	114.44	110.54	175.01	1.04:1:1.58
	Female	25.83	14.43	59.74	37.64	19.74	42.62	32.28	23.80	43.93	32.05	26.26	41.67	127.80	84.23	187.96	1.52:1:2.23
90 day	Male	30.22	18.79	50.98	31.35	46.37	22.26	24.48	25.43	49.76	27.73	24.35	47.92	114.12	112.94	170.94	1.01:1:1.51
	Female	31.91	16.24	52.67	32.14	20.63	42.23	24.19	27.56	48.25	28.66	22.37	48.95	116.90	86.80	192.10	1.35:1:2.21

#### DISCUSSION

Achatina fulica being herbivorous pulmonate gastropod, the lipid classes and fatty acids in this snail are probably exogenous in origin as true in the case of certain bivalves (Beninger & Stephan, 1985). The snails starve during aestivation and during this period females (?) make preparatory steps for future egg laying. It is evident that total lipid profiles of all the organs of A. fulica show distinct male/female differentiation. Total lipid of the digestive gland in both male/female groups reduces up to 30-45 days of aestivation after which it stabilizes in female while it increases in male, which is indicative of more lipid utilization in female than male during aestivation. In a study conducted without sex bias, Mitra and Sur (1989) record decrease of total and neutral lipid in A. fulica from 60 days of aestivation. Higher amount of lipid in digestive gland is reported in active snails and bivalves (Wenne & Polak, 1989; Napolitano & Ackman, 1992; Da Silva & Zancan, 1994; Rakshit et al., 1997). Lower amount of lipid in albumen gland, compared to digestive gland can be explained by the fact that the former acts as a reservoir of carbohydrate and protein (Chaki, 1987). A stable level of lipid in the albumen gland is indicative of lipid transportation from digestive gland which is reported in case of pectinid bivalves as important in reproduction (Vassallo, 1973; Robinson et al., 1981; Pazos et al., 1997). Lipid profiles of mantle and foot exhibit initial decline and then maintains stability throughout aestivation indicating the relative non-involvement of these organs in egg production. It was found that the activities of metabolic enzymes during aestivation were higher in digestive gland than of the mantle (Chaki et al., 2008).

In animals phospholipids serve as a membrane component and play a major role in signal transduction. Phospholipids also participate in the transport of triacylglycerol and cholesterol in animals (Conn *et al.*, 2003). The amount of phospholipid gradually decreases to  $1/4^{\text{th}}$  at 90 days of aestivation in both the sexes. Albumen gland of male exhibits higher amounts of phospholipids than female, but the female albumen gland shows less fluctuation and maintains more or less the same amount of phospholipid during aestivation. This indicates the involvement of this lipid class in egg production, which plays a dynamic role during aestivation. As an alternative explanation, it could be that a small proportion of the highly unsaturated fatty acids are converted to prostacyclins or prostaglandins (steroid hormones) to function in directing the assembly of the ovum. The conversion to steroid hormones is probably achieved by the hydrolysis of phospholipids into free fatty acids whenever required. Cytomorphological studies on the ovotestis of A. fulica show well developed gonads (Rakshit et al., 2005), which are attributed to the availability of steroid hormones Seasonal variations in phospholipid are documented in a terrestrial pulmonate gastropod by (Da Silva & Zancan, 1994). Higher amount of phospholipid in mantle and foot and steady decrease of phospholipid in mantle and foot of both sexes is probably due to maintenance of homeostasis and membrane fluidity (Alexandrov, 1977; Cossins & Prosser, 1978). Phytophagous snails contain cholesterol as the major component of sterols (Idler & Wiseman, 1972). In the present experiment the digestive gland of active male A. fulica shows higher amount of cholesterol than female and

this trend is maintained throughout aestivation. Cholesterol shows higher levels in active male snails and may be associated with spermatogenesis as reported in case of bivalves (Napolitano et al., 1993). The results on the fluctuation level can be due to cessation of supply from the digestive gland at later stages of aestivation. Mantle being the membranous part interacts more with the environment and exhibits greater fluctuation in cholesterol which may be attributed to influence of environment as in the case of bivalve Macoma sp. (Jarjebski & Wenne, 1990), however, these results need further clarification. Triacylglycerol (TG) plays an important role in reproductive strategy of mollusks (Napolitano et al., 1992). Triacylglycerols in digestive gland of male and female A. fulica is mostly above 10% level in both active and aestivated stages, the level being higher in males than female although the pattern of fluctuation appears same. In albumen gland of both sexes TG level is lower compared to digestive gland but no difference in pattern of fluctuation was noted. Mantle and foot contain least amounts of TG. The results are in accordance with the findings of Wenne & Polak (1992). To summarize, the highest amount of TG is found in digestive gland, which is probably because this gland appears to be the most active organ during aestivation. Lustrino et al. (2010) reported that triglycerides metabolism in A. fulica is more influenced by photoperiod than cholesterol metabolism. It is interesting to observe that some organs viz., digestive gland, albumen gland and mantle of both sexes show increment of certain lipid classes. This may be due to lipid transportation from a compensating organ or lipid biosynthesis by that organ. There is no existing information regarding existence of such compensating organ. According to previous workers (Hegg, 1977; Mitra & Sur, 1989) lipid and glycogen metabolism provide energy during aestivation in gastropod Pila globosa. Biosynthesis of lipids in active mollusks has been established but in aestivating snails information is scanty. Studies on metabolic adaptation in aestivating Achatina achatina by Umezurike & Iheanacho (1983) revealed that in short-spell forced starved and aestivating snails glycogen degradation and lipid mobilization takes place in foot, digestive gland and heart muscle at the beginning of aestivation. Such mechanism also takes place in other pulmonate (Van der Horst, 1974; Chaki et al., 2008). The above statement can explain depletion of TG in A. fulica up to 15 days of aestivation, but further studies need to be conducted to explain the surge in later stages of aestivation. Basic types of fatty acids occurring in mollusks include C14, C16, C17, C18, C20, C22 and C24 (Misra et al., 2002; Kraffe et al., 2008). Palmitic (16:0), stearic (18:0) and eicosanoic (20:0) acids are predominant saturated fatty acids in terrestrial snails (Voogt 1972; Van der Horst & Zandee 1973; Catalan et al., 1977), freshwater (Dembitsky et al., 1994) and marine (Saito & Hashimoto, 2010) snails. Fluctuation in saturated fatty acids (SFA) is noted in pulmonate A. fulica, which can be attributed to the fact SFA serves as energy source in bivalves and gastropods (Polak et al., 1987). Variation in 14:0 and 16:0 acids in A. fulica in active as well as aestivating stages can be explained as the result of *de novo* synthesis which occurs in bivalves (Sprecher & James, 1979). Higher percent of 16:0 and 18:0 acids in albumen gland of female

throughout aestivation is possibly related to vitellogenesis. It appears interesting that 18:0 is the predominating fatty acid in digestive gland while 16:0 is predominating in other organs. Sum of 16:0 acids of the four organs of the active snail shows higher value than 18:0 acids, which can be explained by Ackman's (1982) observations which inferred that 16:0 is the principal fatty acid at all evolutionary and trophic levels. This is true in case of freshwater gastropods (Dembitsky et al., 1993b, c, 1994; Misra et al., 2002). Monoenoic constitutes the least amount of fatty acid classes in all the organs of both sexes. Increase in 18:1 acid probably is the reason for increase in monoenoics in digestive gland in male during midaestivation stages. This fact is contradictory to Ackman (1980). In female 18:1 appears to be predominant in contrast to 16:1, which is undetectable after 7<sup>th</sup> day of aestivation. In male albumen gland, 16:1 and 18:1 are the major monoenoics, while in female 18:1 appears to be the chief one and 16:1 drops down at 30<sup>th</sup> day and is undetectable in the later stages. Most probably 18:1 plays a major role in vitellogenesis. Ackman (1980) provided several plausible methods of bioconversion of fatty acids in mollusks, of which  $16:1\omega7 + 2C \rightarrow 18:1\omega7$  seems the most appropriate mechanism in this gastropod. Similar bioconversion pathways are noted in active freshwater snails (Dembitsky et al., 1993b, c; 1994). All these factors point to a clear male/female distinction from the point of bioconversion and utilization of monoethylenic acids (Tables 5-8, 10). Polyunsaturated fatty acids (PUFA), particularly  $C_{18}$  and  $C_{20}$  contribute a major amount of fatty acids. PUFAs show maximum amount in digestive gland, lowest in albumen gland and similar amounts in mantle and foot. Predominating polyenes in active and aestivating stages include 18:2w3 and 20:3w3 acids in varying amounts. Most of these acids can be assumed to be dietary in origin as noted in bivalves (Ackman 1982). Arakelova et al. (2009) are of the opinion that the occurrence and fluctuation of  $\omega$ 3 fatty acids is contingent upon the habitat and motor activity of sea and freshwater mollusks. Significant amount of 20:503 acid is observed in foot of female and albumen gland of male while lesser amount is recorded in albumen gland of female and this can be attributed to bioconversion or biosynthesis. Thus it can be inferred that digestive gland acts as the source of free fatty acids, which can be correlated, to trophic metabolic function of midgut (digestive) gland (Catalan et al., 1977). Low level of 20:306 acid is observed in female digestive gland from 7th-60th day of aestivation while higher amounts are recorded in albumen gland of the same. This is contrary to the levels of  $18:3\omega 3$  acid in female. This is a strong indication of bioconversion through chain elongation process. Steady levels of 18:2w6 acids in mantle of aestivating snails of both sexes can be attributed to an adaptation for minimization of water loss (Wertz & Downing 1990). Linolenic acid  $(18:3\omega 6)$  is probably the precursor for PUFA synthesis in this gastropod. Considerable amounts of 20:4 acid is detected in both digestive and albumen glands in both sexes which is probably dietary in origin. According to Sepe et al. (1998), 20:4 is of dietary origin in edible bivalves and indicator of salinity in marine bivalves (Wenne & Polak, 1992). Occurrence of 20:3\omega3 acid appears to be high in this

pulmonate as compared to marine bivalves (Napolitano et al., 1992) and freshwater gastropods (Misra et al., 2002), but their exact role needs to be ascertained. Thus, total amount of fatty acids in all four organs of A. fulica shows that SFA in male is higher than that of the female up to 45<sup>th</sup> day of aestivation; monoenoics are lower than female in active snails but higher in aestivating stages (Table 10). Polyenes are always higher in female than male except the 15<sup>th</sup> day of aestivation. The fluctuation in the levels of polyenes can be explained due to decreased biological activity of membrane-related processes which occur in conjunction with the reduction of mitochondrial aerobic metabolism observed during aestivation (Stuart et al., 1998). The fatty acid data clearly indicate male/female dimorphism, and also emphasizes monoene to polyene conversion in this gastropod. Monoene: polyene ratio is higher in females after 15<sup>th</sup> day of aestivation whereas higher amounts of SFA in females were observed in contrast to males where it got depleted (Table 10). It is an interesting finding that female snails derive more SFA through conversion from glycogen (neolipogenesis) to maintain supply of polyenes, which is important for vitellogenesis Ratio of SFA: MFA: PUFA (Table 10) supports this hypothesis. According to previous workers (George and Desai, 1954; Krishnamoorthy 1968; Swami & Reddy, 1978; Mitra & Sur, 1989) glycogen depletion during aestivation is related with energy requirement (cf. Chaki et al., 2008) as against lipid metabolism. Thus, it can be stated that glycogen depletion is also important for fatty acid conversion for vitellogenesis. Ovotestis requires more enzymatic activities in metabolic front for gametogenesis during aestivation over other organs (Chaki et al., 2008). This fatty acid pool probably comes through glycogenolysis in the digestive gland. It is to be mentioned that all the above workers carried out the studies without male/female bias.

It can thus be inferred that, the digestive gland is the major lipid storage site and also plays important role in transportation of lipid to other organs during aestivation and maintains higher fatty acid percent than other organs. The other organs studied in this snail maintain a lipid pool. During aestivation, males utilize less amount of lipid for physiological maintenance as compared to females which for vitellogenic utilize it purposes. Phospholipid/cholesterol ratio is always higher in digestive gland and albumen gland of active and aestivating females. In aestivating female, monoenes of digestive and albumen glands are depleted to vield more polyenes, a phenomenon probably required for egg formation. This observation is also supported by higher SFA: MFA: PUFA ratio in female. Overall depletion in fatty acid level during aestivation in both male and female is due to slow and gradual use of these components towards maintenance of the minimum physiological demand. Mantle and foot of both sexes show similar and little variation in lipid and fatty acid storage pattern. Therefore, it is probable that the major site of bioconversion of lipids is digestive gland and there is clear indication of a sexual dimorphism from the point of lipid requirement and metabolism in this hermaphrodite snail. That the hypothesis of the present work that a male/female functional distinction in hermaphrodite snail exists.

#### ACKNOWLEDGEMENT

The authors are thankful toHead, Department of Organic Chemistry, University of Calcutta, 92 A. P. C. Road, Kolkata 700 009, India for providing the necessary laboratory facilities and for extending help towards GLC analysis.

#### REFERENCES

Abad, R., Martinez, D., Mosquera, G. & Sanchez, J.L. (1995) Seasonal variations of lipid classes and fatty acids in flat oyster *Ostrea edulis* from San Cibran (Galicia, Spain). Comp. Biochem. Physiol., 110 C: 109-118.

Ackman, R.G. (1982) Fatty acid metabolism in bivalves. In: Proc. 2nd Intern. Conf. Aquaculture Nutr: Biochem. Physiol. Approaches to Shellfish Nutr., Pruder G D, Llandudno CJ & Conklin DE (eds.) pp. 358-376Louisiana State Univ., Special Publ. No. 2, Baton Rouge, LA, 1982..

Alexandrov, V.A. (1977) Saturation of fatty acids and the temperature conditions of life. In: *Cells, Molecules and Temperature*. Springer, Berlin. pp. 243-255.

Arakelova, E.S., Chebotareva, M.A., Zabelinski, S.A. & Ivanova, V.P. (2009) About effect of habitat and motor activity of molluscs on fatty acid composition of triglycerides and phospholipids (In Russian). Zh Evol Biokhim Fiziol.. 45: 46-52.

Barber, B.J. & Blake, N.J. (1981) Energy Storage and Utilisation in relation to gametogenesis in *Argopecten irradians concentricus* (Say.)J. exp. mar. Biol. Ecol., 52: 121-134.

Beninger, P.G. & Stephan, C. (1985) Seasonal variations in the fatty acids of the triacylglycerols and phospholipids of two populations of adult clam (*Tapes deccusatus* L. and *T. phillipinarum*) reared in a common habitat. Comp. Biochem. Physiol. 81B : 591-601.

Bligh, E.G. & Dyer, W.J. (1959) A rapid method of total lipid extraction and purification. Can. J. Biochem. Physiol., 37: 911-917.

Bradshaw, S.A., O'Hara, S.C.M., Corner, E.D.S. & Eglington, G. (1991) Effects on dietary lipids on the marine bivalve *Scrobicularia plana* feeding in different modes. J. Mar. biol. Assoc. U.K., 71: 635-653.

Catalan, R.E., Castillon, M.P. & Rallo, A. (1977) Lipid metabolism during development of the mollusc *Arion empiricorum*. Distributionof lipids in midgut gland, genitalia and foot muscle. Comp. Biochem. Physiol., 57B: 73-79.

Chaki, K.K. (1987) Some biochemical and histochemical observations on active and aestivating pestiferous mollusc, *Achatina fulica*. Ph.D. Thesis. University of Calcutta, Calcutta.

Chaki, K.K., Dutta, C., Dasgupta, J., Sarkar, A.K. & Misra, K.K. (2008) Profile of some etabolic enzymes of *Achatina fulica* (Pulmonata: Gastropoda) during aestivation. Proc. zool. Soc., Calcutta, 61: 39-50.

Chen, P.S., Toribare, T.Y. & Warner, H. (1956) Microdetermination of phosphorus. Anal. Chem., 28: 1756. Chétail, M. & Fournié, J. (1986) Histological study of the post-embryonic evolution of the hermaphrodite gland from hatching to death in *Derocera reticulatum* (Gastropoda: Pulmonata). Proceedings of the 8<sup>th</sup> International Malacological Congress, Budapest, 1983. In Pintér, L. ed pp. 53 – 58.c/o Hungarian Natural History Museum, Budapest, Hungary.

Conaway, C.A., Fried, B. & Sherma, J. (1996) Effects of restricted food intake on neutral lipid and free fatty acid levels in the faces of *Biomphalaria glabrata* (Gastropoda). Biomed. Chromatogr., 10: 186-188.

Conn, E.E., Stumpf, P.K., Brueni, & G., Doi, R.H. (2003) Outlines of Biochemistry. 5<sup>th</sup> Ed., John Wiley, Singapore.

Cossins, A.R. & Prosser, C.L. (1978) Evolutionary adaptation of membranes to temperature. Proc. natn. Acad. Sci., U.S.A., 75: 2040-2043.

Cunha, R., Rodrigues, A. & Sousa, H. (1998) The albumen gland as a maturation diagnostic organ in *Oxychilus* (*Drouetia*) atlanticus (Stylommatophora, Zontidae). Proc. 8th Int. Cong Inv. Repr. Dev., Amsterdam, 1998.

Da Silva, R.S.M. & Zancan, D.M. (1994) Seasonal variation of the carbohydrate and lipid metabolism in a land pulmonate gastropod, *Megalobulimus oblongus*. Comp. Biochem. Physiol., 108A : 337-341.

Dembitsky, V.M., Kashin, A.G. & Stefanov, K. (1992) Comparative investigation of phospholipids and fatty acids of freshwater molluscs from the Volga river basin. Comp. Biochem. Physiol., 102B: 193-198.

Dembitsky, V.M., Kashin, A.G. & Stefanov, K. (1993a) Comparative examination of phospholipids and fatty acids from some Caspian invertebrates. Comp. Biochem. Physiol., 104B: 617-622.

Dembitsky, V.M., Rezanka, T. & Kashin, A.G. (1993c) Comparative study of the endemic freshwater fauna of Lake Baikal - I. Phospholipid and fatty acid composition of two mollusc species, *Baicalia oviformus* and *Benedicta baicalensis*. Comp. Biochem. Physiol., 106B: 819-823.

Dembitsky, V.M., Rezanka, T. & Kashin, A.G. (1994) Comparative study of the endemic freshwater fauna of lake Baikal - IV. Phospholipid and fatty acid compositions of two gastropod molluscs of genus *Valvata*. Comp. Biochem. Physiol., 107B: 325-330.

Dembitsky, V.M., Rezanka, T. & Kashin, A.G. (1993b) Fatty acid and Phospholipid composition of freshwater molluscs *Anadonta piscinalis* and *Limnaea fragilis* from the river Volga. Comp. Biochem. Physiol., 105B: 597-601.

Dutta, A., Pal, P.K., Ghosh, A., Misra, S., Nandi, S. & Choudhury, A. (1986) Lipids and fatty acids of the gastropod mollusc *Cerethidea cingulata*. J. Am. Oil. Chem. Soc., 63: 223-225.

Egonmwan, R. I. (2007) Gross Morphology and Ultrastructural Study of Albumen Gland of the Land Snail *Archachatina marginata ovum* (Pfeiffer) (Pulmonata: Achatinidae). Pak. J. Biol. Sci. 10: 322-325.

Ekin, I. (2012) Variations of fatty acid contents in selected tissues of freshwater mussel *Corbicula fluminalis* (Mollusca: Bivalvia) from Tigris River. Energy Education Science and Technology Part A: Energy Science and Research, 29: 157-164.

Ekin, I., Ba han, M. & e en, R. (2012) Fatty acid composition of lipid classes in two mussel populations (*Anodonta piscinalis* and *Corbicula fluminalis*) living in Tigris River. Turk J Biochem, 37: 283–289.

Frazer, B.A., Reddy, A., Fried, B. & Sherma, J. (1997) HPTLC determination of neutral lipids and phospholipids in *Lymnaea elodes* (Gastropoda). J. Planar Chrom. Modern TLC. 10: 128-130.

Fried, B., Frazer, B.A., Lee, M.S. & Sherma, J. (1998) Thinlayer chromatography and histochemistry analyses of neutral lipids in Helisoma trivolvis and of larval trematodes. Parasitol Res., 84: 369-373.

Gallager, S.M. & Mann, R. (1986) Growth and survival of larvae of *Mercenaria mercenaria* (L.) and *Crassostrea virginica* (Gmelin) relative to brood stock conditioning and lipid content of eggs. Aquaculture, 56: 105-121.

George, J.C. & Desai, B.N. (1954) On the liver fat in *Pila* globosa (Swainson). J. Anim. Morph. Physiol., 1: 56-58.

Ghosh Choudhuri, P., Chakraborty, M.M. & Bhattacharyya, D.K. (1983) Modification of some tree-borne seed fats for the preparation of high priced confectionary fats. Fette Seifen Aust., 85: 224-229.

Gordon DT & Collins N 1982. Anatomical distribution of sterols in oysters (*Crassostrea gigas*). Lipids, 17: 811-817.

Hegg, J. (1977) Oxygen consumption and the use of metabolic reserves during starvation and aestivation in *Bulinus (Physopsis) africanus* (Pulmonata: Planorbidae). Malacologia, 16: 549-560.

Hyman, L. H. (1967) The Invertebrates. Vol. 6 Mollusca. I. McGraw-Hill, New York.

Jarzebski, A. & Wenne, R. (1990) Sterol composition of marine bivalve from the Genus *Macoma*. Comp. Biochem. Physiol. 97B: 81-82.

Jarzebski, A., Wenne, R. & Habermehl, G. (1986) Anatomical distribution of lipids and sterols in *Macoma balthica*. Comp. Biochem. Physiol. 85B: 135-137.

Kates M 1972. Techniques in Lipidology. American Elsevier Publs. New York

Klingensmith, J.S. & Stillway, L.W. (1982) Lipid composition of selected tissues of the hardshell clam, *Mercenaria mercenaria*. Comp. Biochem. Physiol. 71B:111-112.

Kluytmans, J.H., Boot, J.H., Oudejans, R.C.M. & Zandee, D.J. (1985) Fatty acid synthesis in relation to gametogenesis in the mussel *Mytilus edulis* L. Comp. Biochem. Physiol. 81B: 959-963.

Kraffe, E., Grall, J., Le Duff, M., Soudant, P. & Marty, Y. (2008) A striking parallel between cardiolipin fatty acid composition and phylogenetic belonging in marine bivalves: a possible adaptative evolution? Lipids, 43: 961-970.

Kraffe, E., Soudant, P., Marty, Y. & Kervarec, N. (2005) Docosahexaenoic acid- and eicosapentaenoic acid-enriched cardiolipin in the Manila clam Ruditapes philippinarum. Lipids, 40: 619-625.

Kraffe, E., Soudant, P. & Marty, Y. (2004) Fatty acids of serine, ethanolamine, and choline plasmalogens in some marine bivalves. Lipids, 39: 59-66.

Krishnamoorthy, R.V. (1968) Hepatopancreatic unsaturated fatty acids during aestivation of the snail, *Pila globosa*. Comp. Biochem. Physiol. 24: 279-282.

Luchtel, D.L. (1972) Gonadal development and sex determination in pulmonate molluscs. 1. *Arion circumscriptus*. Zeiscrift für Zellforschung, 130: 279-301.

Lustrino, D., Tunholi-Alves, V.M., Tunholi, V.M., Marassi, M.P. & Pinheiro, J. (2010) Lipids analysis in hemolymph of African giant *Achatina fulica* (Bowdich, 1822) exposed to different photoperiods. Braz J. Biol. 70: 129-34.

Mangold, H.K. (1969) Thin layer chromatography. Stahl. E. (ed.) Springer Verlag, New York, 263-421.

Misra, K.K., Shkrob, I., Rakshit, S. & Dembitsky, V.M. (2002) Variability in fatty acids and fatty aldehydes in different organs of two prosobranch gastropod mollusks. Biochem. Syst. Ecol, 30: 749-761.

Misra, S., Misra, S., Choudhury, A. & Ghosh, A. (1986) Lipids and fatty acids of the gastropod mollusc, *Cerethidea obtusa*. Food Chem., 22: 251-258.

Mitra, S. & Sur, R. (1989) Changes in the lipid and carbohydrate contents of the digestive gland during aestivation of two gastropods *Achatina fulica* and *Pila globosa*. Environ. Ecol. 7: 658-662.

Mukai, S.T., Hpque, T. & Morishita, F. (2004) Cloning and characterization of a candidate nutritive glycoprotein from the albumen gland of the freshwater snail *Heliosoma duryi* (Mollusca:Pulmonata). Invert. Biol. 123: 63-92.

Napolitano, G.E. & Ackman, R.G. (1992) Anatomical distributions and temporal variations of lipid classes in sea scallops *Placopecten magellanicus* (Gmelin) from Georges Bank (Nova Scotia). Comp. Biochem. Physiol. 103B: 645-650.

Parivar, K. (1980) Differentiation of the Sertoli cells and post reproductive epithelial cells in the hermaphrodite gland of *Arion ater* (L.) (Mollusca, Pulmonata). J. Moll. Stud. 46: 139-147.

Pazos, A.J., Roman, G., Acosta, C.P., Sanchez, J.L. & Abad, M. (1997) Lipid classes and fatty acid composition in the female gonad of *Pecten maximus* in relation to reproductive cycle and environmental variables. Comp. Biochem. Physiol., B117: 393-402. Polak, L., Jarzebski, A., Wenne, R. & Falkowski, L. (1987) Seasonal changes in condition and lipid composition of the bivalve *Macoma balthica* L. from the Gulf of Gdansk (Southern Baltic). Comp. Biochem. Physiol. 88B: 881-885.

Rakshit, S., Bhattacharyya, D.K. & Misra, K.K. (1997) Distribution of major lipids and fatty acids of an estuarine gastropod molluscs *Telescopium telescopium*. Folia Biol. (Krakow), 45: 83-87.

Rakshit, S., Chaki, K.K., Sarkar, A.K. & Misra, K.K. (2005) Cytomorphology of ovotestis of a pulmonate snail, *Achatina fulica*. Proc. zool. Soc., Calcutta, 58: 93-104.

Robinson, W.E., Wehling, W.E., Morse, P.M. & MeLeod, G.C. (1981) Seasonal changes in soft-body component indices and energy reserves in the Atlantic deep-sea scallop *Placopecten magellanicus*. Fish. Bull. 79: 449-458.

Saito, H. & Hashimoto, J. (2010) Characteristics of the fatty acid composition of a deep-sea vent gastropod, *Ifremeria nautilei*. Lipids, 45: 537-48

Sepe, N., De Petrocellis, L., Montanaro, F., Cimino, G. & Di Marzo, V. (1998) Bioactive long chain N-acylethanolamines in five species of edible bivalve molluscs and its role in physiology and sea food industry. Biochim. Biophys. Acta, 1389: 101-111.

South, A. (1992) Terrestrial Slugs, Chapman and Hall, London

Sprecher, H. & James, A.T. (1979) Biosynthesis of long chain fatty acids in mammalian systems. In: Geometrical and Positional Fatty Acid Isomers. Emken, E. A. and Dutton, H. J.eds. pp. 303-338.American Oil Chemists' Society, Champaign, IL.

Stoilov, I., Papov, S., Marekov, N. & Andreev, St. (1984) Anatomical distribution of sterols in some filter feeding invertebrates. Comp. Biochem. Physiol. 79B: 225-228.

Stuart, J.A., Gillis, T.E. & Ballantyne, J.S. (1998) Remodeling of phospholipid fatty acids in mitochondrial membranes of estivating snails. Lipids, 33: 787-93.

Swamy, K.S. & Reddy, Y.S. (1978) Adaptive changes to survival during aestivation in the gastropod snail, *Pila globosa*. J. Sci. Ind. Res. 37: 144-157.

Swift, M.L., White, D. & Ghassemieh, M.G. (1980) Distribution of neutral lipids in the tissues of the oyster *Crassostrea virginica*. Lipids, 15: 129-132.

Tomiyama, K. (1995) Mate choice in a simultaneously hermaphroditic land snail, *Achatina fulica* (Stylommatophora, Achatinidae). Proc. 12th Intern.Malac. Cong., Vigo (1995), 279-280 (Abs.).

Umezurike, G.M. & Iheanacho, E.N. (1983) Metabolic adaptations in aestivating giant African snail *Achatina achatina*. Comp. Biochem. Physiol. 74B: 493-498.

Vale, P. (2010) Profiles of fatty acids and 7-O-acyl okadaic acid esters in bivalves: can bacteria be involved in acyl esterification of okadaic acid? *Comp Biochem* Physiol C Toxicol Pharmacol. 151: 18-24.

Van der Horst, D.J., Kingma, F.J. & Oudejans, RCHM (1973) Phospholipids of the pulmonate land snail *Cepaea nemoralis* (L.). Lipids, 8: 759-765.

Van der Horst, D.J. & Oudejans, RCHM (1976) Fate of dietary linoleic and linolenic acids in the land snail *Cepaea nemoralis* (L.). Comp. Biochem. Physiol. 55B: 167-170.

Van der Horst, D.J. & Zandee, D.I. (1973) Invariability of the composition of the fatty acids and other lipids in the pulmonate land snail *Cepaea nemoralis* (L.) during an annual cycle. J. comp. Physiol. 85: 317- 326.

Van Handel, E., Zilversmit, D.B. & Tenn, M. (1957) Micromethod for the direct determination of serum triglyceride. J. Lab. Clin. Med. 50: 152.

Vassallo, M.T. (1973) Lipid storage and transfer in the scallop, *Chlamys hericia* Gould. Comp. Biochem. Physiol. 44A: 1169-1175.

Viarengo, A., Accomando, R., Roma, G., Benatti, U., Damonte, G. & Orunesu, M. (1994) Differences in lipid composition of cell membranes from Antarctic and Mediterranean scallops. Comp. Biochem. Physiol. 109B: 579-584.

Voogt, P.A. (1972) Lipid and sterol components and metabolism in mollusca. In: Chemical Zoology, Vol. 7. Florkin, M. and Scheer, B. T. eds. pp. 329-370 New York and London.

Wenne, R. & Polak, L. (1989) Lipid composition and storage in the tissues of the bivalve *Macoma balthica*. Biochem. Syst. Ecol. 17: 583-587.

Wenne, R. & Polak, L. (1992) Lipid storage strategy in the clam *Macoma* (Bivalvia). Proc. 12th Baltic mar. Biol. Symposium, pp. 179-182.

Wertz, P.W. & Downing, D.T. (1990) Metabolism of linoleic acid in porcine epidermis. J. Lipid Res. 31: 1839-1844.

Zandee, D.I., Holwerda, D.A. & Zwaan, de A. (1980) Energy metabolism in bivalves and cephalopods. In: *Animals and Environment Fitness*. Gilles, R. ed, pp. 185-206, Pergamon, Oxford.

Zlatkis, A., Zak, B. & Boyle, A. (1953) A method for the determination of serum cholesterol. J. Lab. clin. Med. 41: 486-490