



IMPACT OF INDUCED MUTATIONS ON THE GROWTH PERFORMANCE AND YIELD OF S₁₃ MULBERRY VARIETY USING GAMMA-RAYS

¹Muniswamy Reddy, P.M. & ²Munirajappa

Central Tasar Research and Training Institute, (Central Silk Board, Ministry of Textiles) Piska Nagri Ranchi -835303, Jharkhand, India,
Moriculture Laboratory, Department of Studies in Sericulture, Bangalore University, Bangalore 560 056, India

ABSTRACT

S₁₃ is a drought tolerant mulberry variety cultivated under rainfed condition. In order to increase the leaf yield and to improve the quality the hard wood stem cuttings of this taxon were irradiated with γ -rays (4kR, 5kR, 6kR, 7kR, 8kR, 9kR and 10kR). Growth attributes such as sprouting performance, rooting ability, survivability, yield and phytochemical components were recorded in the irradiated population. Comparable controls were also maintained. It is observed that the lower dosages of gamma - rays i.e., 4kR and 5kR were less effective in inducing variability. At higher dosages viz., 8kR, 9kR and 10kR deformities like delayed sprouting, poor rooting, weathering of inflorescence, stunted growth, plants with weak and feeble branches, plants bearing small leaves with wrinkled and coriaceous texture were observed. On the other hand, at 6kR beneficial mutants were recorded. These variants exhibited superiority over the control in yield, phytochemical constituents and commercial characteristic features of the cocoon. The present study recommends the exploitation of these promising mutants of S₁₃ for commercial exploitation.

Key words: Gamma irradiation, Diploid, S₁₃ mulberry, Chemo-assay, Bioassay.

INTRODUCTION

Sericulture is one of the major agro-industry that plays an important role in uplifting rural economy and provides employment opportunity for rural folk. The silkworm *Bombyx mori* L. is a monophagous insect, cherishing on its sole food plant i.e., mulberry. Mulberry leaf is known to be rich in protein, starch and vitamins, helping the silkworm to meet its dietary requirements. Mutation breeding employing radiation (X-rays, Gamma rays, Fast neutrons, etc.) was initiated to explore the feasibility of developing new varieties. Mutagenesis has been proved to be extremely useful in creating a new variability in the existing gene pool. Of late, genetically effective radiations have been widely used for induction of mutation in mulberry (Hazama, 1967, 1968a, 1968b; Katagiri and Wada, 1971; Nakajima, 1973; Fujita, 1974; Kukimura, *et al.*, 1975, 1976; Aliev, 1977; Fujita, *et al.*, 1980; Jayaramaiah and Munirajappa, 1987; Ramesh, 1997). Gamma - rays induced diploids have shown all the symptoms such as larger leaf size, bigger leaf area, dark green colored leaves, delayed flowering, slower growth etc., when compared to control. Keeping the above aspects in view, the present investigation aims at improving the already existing cultivars by using gamma irradiation.

MATERIALS AND METHODS

Hardwood stem cuttings of S₁₃ mulberry genotypes were selected from young and healthy bushes. Each cutting measuring about 6-8 inches in length half inch in diameter bearing 3-4 active vegetative buds was preferred cuttings were irradiated with 4kR 5kR, 6kR, 7kR, 8kR, 9kR and 10kR doses of gamma rays from a Co 60 gamma rays source at Kidwai Memorial Institute of Oncology Bangalore-560021. In each dose 20 cuttings in three

replications were used. The irradiated cuttings were planted in polythene bags and maintained providing necessary agricultural inputs for their further response. Comparable controls (untreated cuttings) were also maintained. Growth and Morphological characters of saplings were observed included sprouting, rooting, survivability percentage, plant height, number of branches, length of the petiole, leaf area etc., (R₁ generation). Phytochemical constituents and commercial characteristic features of cocoons were analyzed by following standard procedures (R₂ generations).

RESULTS

The stem cuttings of this taxon irradiated with different doses (4kR to 10kR) of gamma rays showed varied responses in sprouting behaviour. In general, sprouting percentage ranged from 45.00% to 76.66%. Similarly, number of days taken for sprouting differed with different doses of radiation. In control, the cuttings took 9 days for sprouting with a sprouting percentage of 86.66%. In the present study, it is evident that root development, decreased in a linear trend with the increase of gamma irradiation dosage. Control plants showed 85.00% rooting which is the optimum value required for the commercial exploitation of mulberry genotypes. Rooting percentage varied from 80.00% to 40.00% in different doses of gamma irradiation. Cuttings irradiated with 4kR and 5kR gamma rays showed slightly decreased rooting ability when compared to control. Further, cuttings, which are administered with 6kR gamma rays revealed better rooting ability (85.00%). Saplings recovered from the cuttings irradiated at 6kR gamma rays showed extensive root system both in spread and depth when compared to control. In the irradiated populations, survival percentage

Impact of gamma-rays on the growth performance and yield of S₁₃ mulberry variety

ranged from 68.33% to 43.33% when compared to control (80.00%). At higher doses (7kR to 10kR), this genotype exhibited poor growth with feeble and weak branches. On the other hand, reduction in survivability was not significant at lower doses of gamma rays (4kR to 6kR). Survival percentage was maximum at 6kR (73.33%) and minimum at 10kR (43.33%). LD₅₀ for this taxon was found to be around 8kR since the survivability percentage was 51.66% (Table-1).

Plant height and leaf area was found to be decreased with increase in dose of gamma rays showing negative correlation. The marginal reduction in plant height, stunted growth, deformities in growth like fasciation, plant with weak and feeble branches, plants bearing small leaves with wrinkled and coriaceous texture were noticed in the populations of cuttings irradiated at 9kR and 10kR. However, some of the populations irradiated with 6kR gamma rays exhibited increased plant height when compared to control. These variants also showed increasing leaf area with thick, succulent and dark green leaves (Fig-1). Significant results were noticed with respect to internodal distance, many irradiated saplings revealed decreased internodal distance when compared to control (Table-1).

No significant variation was observed in length of the inflorescence at any of the treatments. Inflorescence at higher dosages did not attain maturity, often-deformed inflorescence was observed. Pollen fertility lowered which gradually decreased with increased dosage of gamma rays. Total foliage yield per plant and average weight of leaves (100Nos) was greater (305.21g) when compared to control (268.27g). Initially, the average yield determined to be 0.645kg/bush compared to 0.535kg in the control counterparts in the R₂ generation.

PHYTOCHEMICAL STUDIES

Protein, reducing sugars, amino acids, chlorophyll-a, chlorophyll-b and total chlorophyll were estimated in the mutants grown in the R₂ generation. In the mutant, the amount of proteins in tender, medium and coarse leaves were 25.82%, 24.38% and 24.30% respectively which are higher than the protein percentage in the control (25.38%, 24.11 % and 23.72 %) (Table-2).

Amount of reducing sugar present in tender medium and coarse leaves of induced mutant were 3.45 %, 3.59 % and 3.24 % respectively which are higher than the control (Table-2).

Amino acid content was estimated to be 12.60μ moles/gf.wt., 12.31μ moles/ gf.wt. and 11.84μ moles/gf.wt. respectively in tender, medium and coarse leaves of induced mutants as against the values 12.24μ moles/ gf.wt., 11.90μ moles/gf.wt. and 10.91 μmoles/g f.wt. found in control (Table- 2)

Chlorophyll-a was found to be 2.15, 2.17 and 2.16mg/gf.wt. respectively in tender, medium and coarse leaves of induced mutants compared to 2.04, 2.03 and 2.14mg/gf.wt. for the control. Chlorophyll-b and total chlorophyll were also found to be higher in induced mutants compared to control plants of S₁₃ mulberry variety (Table-2).

Moisture percentage was found to be slightly higher in mutant with 75.47%, 74.51% and 73.38% in tender, medium and coarse leaves respectively as against 75.07%,

74.32% and 72.82% of the control. Similarly, moisture retention capacity at 6 hrs and 12 hrs of mutant was also found to be higher than the control (Table-2).

Various commercial characteristics like ERR, Larval weight, Cocoon weight, Shell weight, Shell weight percentage, Filament length, Reelability percentage, Renditta and Denier were recorded. Larvae fed with the leaves of induced mutants showed an average weight of 3.38gms, which was higher than control (3.26gms). Average cocoon and shell weight were found to be 1.80gms and 0.29gms respectively in case of mutants compared to control (1.67gms and 0.26gms). Higher shell weight percentage was also encountered in the mutants. Similarly filament length was found to be was higher in the mutants (757.94mts) compared to control (685.125mts). Renditta and Denier were also found to be better in the leaves of induced mutants of S₁₃ mulberry variety (Table-3).

DISCUSSION

In the present study, Gamma irradiation studies revealed that the lower dosages (4kR & 5kR), in general were less productive in the induction of mutations whereas at the higher dosages (8kR, 9kR & 10kR) deformity and inhibition of growth leading to semi lethality to complete lethality were observed. The morphological leaf mutants like oblong leaf with wrinkled lamina, variants with slender branches bearing small deformed leaves and mutants with increased lamina (Broad leaves) were secured in irradiated populations. Similar findings have been reported in large number of mulberry genotypes obtained through irradiation by several workers (Katagiri, 1970; Tarar, 1970; Kuchkarov and Ogurtrou, 1987; Dandin *et al.*, 1996) On the other hand, the moderate dosage like 6kR was found to be fruitful in the induction of beneficial mutants. Hazama *et al.*, (1968) studied the varietal defences to radio sensitivity and bud mutations of mulberry trees in gamma irradiation populations and found the inhibition of height and branching pattern. Similar results have also been reported by Rao, *et al.*, (1984), Jayaramaiah (1987) and Ramesh (1997) in S₅, S₅₄ and local mulberry varieties due to the effect of irradiation. Katagiri (1976a) reported deformation in leaf and inhibition of growth at higher doses of gamma rays in mulberry variety Ichinose.

Induced mutants of S₁₃ variety exhibited positive increased trend in the amount of protein, reducing sugar, amino acid and chlorophyll content which is important from the nutritional point of silkworm (Ito and Arni, 1963, Ito and Nimura, 1966). Protein is the main constituent of mulberry leaf which plays a vital role in the development of silk gland (Anfinsen, *et al.*, 1958; Fukuda, *et al.*, 1959; Qader, 1987; Bongale, *et al.*, 1993, 1995; Bose, *et al.*; 1991). It has been proved that the silk worm derives about 70% of proteins from the mulberry leaves for the biosynthesis of silk (Fukuda, *et al.*, 1956; Kaware, 1975). Amino acid and reducing sugar are also required for the synthesis of fibroin and sericin (Vijayan, *et al.*, 1997; Dorcus and Vivekanandan, 1997; Sastry, *et al.*; 1988). Effect of moisture content on silkworm rearing is well documented (Ito, 1963). Low leaf moisture is known to affect the growth and development of silkworm larvae.

Data related to effective rate of rearing, larval weight, cocoon weight, shell weight, shell weight percentage and silk reeled (average filament length, renditta, denier, reelability percentage) were worked out by feeding the leaves of mutants to the cross breed silkworm larvae (PM×NB₄D₂). The mean values for these parameters in regard to both the mutants compared to their respective control counterparts were found to be in the improving trend reflecting the superiority of mutants obtained in the present investigation. The percentage of improvement recorded in the larval weight (6.90%) filament length (8.90%) reelability (7.02%) renditta (7.45%) denier (6.54%) and effective rate of rearing (11.01%) with regard to mutants recovered from S₁₃ mulberry variety. The present investigation clearly establishes the superiority of the induced mutants secured in the present work. Comparable results have also been reported by the earlier workers (Sastry, *et al.*, 1969, Mastfaev, 1968, 1970, 1971; Badalov, 1971; Das and Prasad, 1974; Sikdar, 1993; Bongale and Chaluvachari, 1995; Ramesh, 1997, 2001).

REFERENCES

- Aliev, M.O. (1977) Use of chemical mutagens combined with hybridization of mulberry forms differing in ploidy (in Russian). *Plant Breed. Abstr.*, 49 (1979): No. 5129.
- Anfinsen, J.R., Anson, C.B., Kenneth Bailey, M.L., Edsaii, M.L. and John, T. (1958) Biosynthesis and spinning of silk proteins. *Advances in protein chemistry*, Vol. **13**: 116-118.
- Badalov, N.G. (1971) Result of testing the triploid mulberry Khanalar-tut for silkworm feeding in spring, summer and autumn. *Referativnyi Zhurnal*,
- Bongale, U.D. and Chaluvachari (1995) Evaluation of eight mulberry Germplasm variety by leaf biochemical and bio-assay moulting studies. *Sericologia*, Vol. **35** (1) : 83-94.
- Bongale, U.D., Chaluvachari., Lingaiah., Rao, B.V.N. and Mallikarjunappa, R.S. (1993) Leaf quality evaluation of mulberry gardens in Karnataka. *Trends in Life Sciences (India)*. Vol. **8** (1): 33-37.
- Bose, P. C., Majumdar, S. K. and Sengupta, K. 1991. A comparative biochemical study of six mulberry (*Morus alba* L.) varieties. *Indian J. Seric.*, Vol. **30** (1): 83-87.
- Das, B. C. and Prasad, D. N. (1974) Evaluation of some tetraploid and triploid mulberry varieties through chemical analysis and feeding experiments. *Indian J. Seric.*, Vol. **13**: 17-22.
- Dorcus, D. and Vivekanandan, M. (1997) Exploitation of mulberry genotypes for drought tolerance potential. *J. Seric. Sci. Jpn.*, Vol. **66**(2): 71-80.
- Fujita, H. (1974) Induced dieback disease resistant bud mutations by gamma irradiation in mulberry. *Mutant. Breed. Newsl.*, Vol. **4** : 6.
- Fukuda, T., Sudo, M., Matuda, M., Hayashi, T., Kurose, T., Horiuchi, Y. and Mareen Florrin. (1959) Formation of silk protein during the growth of the silkworm larvae. *Bombyx mori* L. *Proc. 4th Ant. Cong. Biochemistry*, Vol. **12**: 90-112.
- Hazama, K. (1967) On the useful mutants of mulberry induced by radiations. In *Proc. 8th Jpn. Conf. Radioisot.*, pp. 415-417.
- Hazama, K. (1968a) Breeding of mulberry tree. *J.A.R.Q.*, Vol. **3**: 15-19.
- Hazama, K. (1968b) Adaptability of mutant in mulberry tree. *Gamma Field Symp.*, Vol. **7** : 79-85.
- Ito, J.A and Nimura (1966) Referred in: the silkworm; an important laboratory tool. Ed. By Tazima, Y. 1978. Kodensha Ltd., Tokyo.
- Ito, T. and Arai, N. (1963) A food value of mulberry leaves for the silkworm, *Bombyx mori* L. Determined by means of artificial diets. 1. Relationship between kinds of mulberry leaves and larval growth. *Bull. Seric. Expt. Stn. Jpn.*, Vol. **18** 117-120.
- Jayaramaiah, V.C. and Munirajappa (1987) Induction of mutations in mulberry variety 'Mysore Local' by gamma irradiation. *Sericologia*, Vol. **27** (2): 199-204.
- Katagiri, K. (1976a) Radiation damage and induced tetraploidy in mulberry *Morus alba* L. *Environ. Exp. Bot.*, Vol. **16** (2/3): 119-130.
- Katagiri, K. and Wada, M. (1971) Induction of bud mutation resistant to stem blight disease in mulberry by chronic gamma-irradiation. *Jpn J. Breed.*, Vol. **12**: 156.
- Kawase (1975) In: Text book of tropical sericulture. Japan overseas co-operation volunteers, Tokyo: 155.
- Kukimura, H., Ikeda, F., Fujita, H., Maeta, T., Nakajima, K., Nakahira, K. and Somegou, M. 1975, 1976. Genetical, cytological and physiological studies on the induced mutants with special regard to effective methods for obtaining useful mutants in perennial woody plants. I. In: Improvement of vegetatively propagated plants through induced mutations, Tokai, IAEA, Vienna, pp. 83-104.
- Kumar, R. and Mani, S.C. (1997) Chemical mutagenesis in Manhar variety of rice (*Oryza sativa* L.). *Indian J. Genet.*, Vol. **57**: 120-126.
- Machaiah, J.D. and Vakil, U.K. (1979) The effect of gamma in the formulation of α -amylase isoenzyme in germinating wheat. *Envir Exp Bot.*, Vol. **19**: 337-348.
- Mustafaev, A.S. (1970) Biological and economic properties of triploid forms of mulberry. *Polyploidy U. Sheikovitsy*, Moscow: 30-35.
- Mustafaev, A.S. (1968) Study on the effect of leaf quality of experimentally obtained triploid mulberry forms on the

Impact of gamma-rays on the growth performance and yield of s₁₃ mulberry variety biological indices of Chinese silkworm. Doki. Akad. Nank. Az. SSR, Vol. **22** (4): 49-54.

Mustafev, A.S. (1971) Biological and economic properties of triploid form of mulberry. Referativnyi. Zhurnal, Vol. **6** (55): 526.

Nakajima, K. (1973) Induction of useful mutation in mulberry by gamma rays. In induced mutation in vegetatively propagated plants. IAEA, Vienna: 105-117.

Patel, J.D. and Shah, J.J. (1974) Effect of gamma irradiation on seed germination and organization of shoot apex in *Solanum melongena* and *Capsicum annuum* Phytomorphology, Vol. **24**:174-180.

Ramesh, H.L and Munirajappa (2001) Colchicine-induced variability in mulberry variety kosen. Bull. Ind. Acad. Seri., Vol. **5** (2): 34-41.

Ramesh, H.L. (1997) Induction of variability for important morpho-economic traits in mulberry. Ph.D., thesis, Bangalore University, Bangalore.

Reddy, K.J.M and Vidyavathi, (1985) Effect of sumithion on the germination, growth, chromosomal aberration and the enzyme amylase of *Dolichos biflorus* L. J. Indian Bot. Soc., Vol. **64**:88-92.

Sastry, C.R., Venkataramu, C. and Azeez Khan (1969) Induced tetraploid of an improved strain, Kanva-2 of mulberry (*Morus alba* L.) Silkworm. Inf. Bull., Vol. **1** (1): 95-99.

Sastry. C.R., Jolly, M.S., Subramanyan, M.R. and Madhav Rao, Y.R. (1988) Studies on the varietal differences in the loss of moisture from harvested mulberry leaves. Indian J. Seric., Vol. **27**: 85-90.

Selim, A.P., Hussein, H.A.S. and Kishawaf, I I S (1974) EMS and gamma ray induced mutation in *Pisum sativum* L. II Effect of EMS and gamma rays on M₁ generation seedling height and fertility. Egyp J. genet Cytol., Vol. **3** :172-192.

Sikdar, A.K. (1993) Chemo and Bio-assay of high yielding triploids and diploids of mulberry (*Morus* spp.). Indian J. Seric., Vol. **32** (2): 169-174.

Vijayan, K., Raghunath, M.K., Das, K.K., Tikadar, S.P., Chakraborti, B.N. Roy. and Qadri, S.M.H. (1997) Studies on leaf moisture of mulberry germplasm varieties. Indian J. Seric., Vol. **36** (2): 155-157.

TABLE-1. Effect of gamma irradiation on propagation and growth attributes of S₁₃ mulberry variety (R₁ generation)

Treatments	Sprouting (%)	Rooting (%)	Surviv-ability (%)	Height of the plant (cm)	Number of Branches	Internodal distance (cm)	Length of petiole (cm)	Leaf area (cm ²)	Number of flowers/ inflorescence	Length of inflorescence (cm)	Pollen fertility (%)
Control	86.66 ± 1.66	85.00 ± 2.88	80.00 ± 2.88	163.73 ± 2.37	3.66 ± 0.33	3.70 ± 0.01	2.50 ± 0.05	126.14 ± 1.16	39.00 ± 2.64	2.73 ± 0.03	78.41 ± 0.86
4kR	76.66 ± 1.66	80.00 ± 5.77	68.33 ± 1.66	159.46 ± 1.73	4.33 ± 0.33	3.26 ± 0.03	2.66 ± 0.17	135.61 ± 0.87	32.33 ± 2.18	2.62 ± 0.03	72.97 ± 1.29
5kR	71.66 ± 1.66	70.00 ± 2.88	63.33 ± 3.33	155.46 ± 3.18	4.00 ± 0.57	3.63 ± 0.04	2.53 ± 0.14	132.19 ± 0.62	29.66 ± 0.88	2.46 ± 0.07	66.38 ± 3.20
6kR	73.33 ± 3.33	85.00 ± 2.88	73.33 ± 3.33	173.43 ± 3.01	4.66 ± 0.33	3.07 ± 0.06	2.16 ± 0.08	140.58 ± 0.60	31.66 ± 1.20	2.29 ± 0.07	68.00 ± 0.91
7kR	70.00 ± 2.88	65.00 ± 2.88	55.00 ± 2.88	148.16 ± 4.93	4.33 ± 0.88	4.09 ± 0.07	2.90 ± 0.26	129.80 ± 0.32	32.00 ± 1.52	2.42 ± 0.4	61.36 ± 2.17
8kR	60.00 ± 2.88	60.00 ± 2.88	51.66 ± 1.66	133.40 ± 6.32	4.33 ± 0.46	3.97 ± 0.01	3.03 ± 0.12	124.76 ± 1.47	30.33 ± 3.92	2.31 ± 0.06	57.35 ± 3.75
9kR	51.66 ± 4.40	45.00 ± 2.88	45.00 ± 2.88	131.33 ± 5.27	4.00 ± 0.57	4.27 ± 0.09	3.00 ± 0.15	127.30 ± 1.18	25.66 ± 2.60	2.27 ± 0.07	50.68 ± 3.08
10kR	45.00 ± 2.88	40.00 ± 2.88	43.33 ± 3.33	125.80 ± 2.60	3.66 ± 0.33	4.08 ± 0.01	3.06 ± 0.16	120.15 ± 0.46	25.66 ± 2.33	2.20 ± 0.04	44.03 ± 2.43
SEM	3.946	5.088	4.307	5.318	0.709	0.098	0.208	1.379	3.422	0.079	2.898
CD@ 5%	8.464	10.913	9.237	11.407	1.522	0.212	0.447	2.957	7.339	0.171	6.216

TABLE- 2. Phytochemical constituents of S₁₃ mulberry mutant recovered at 6kR gamma irradiation (R₂ generation)

Treatments	Maturity of leaves	Protein (%)	Reducing sugar (%)	Amino acid (μmoles/gf. wt.)	Chl - a (mg/gf.wt)	Chl - b (mg/gf.wt)	Total chlorophyll (mg/gf.wt)	Moisture content (%)	Moisture retention capacity (%)	
									6hrs	12hrs
Control	Tender	25.38 ± 0.17	2.82 ± 0.300	12.24 ± 0.035	2.04 ± 0.005	0.29 ± 0.003	2.33 ± 0.003	75.07 ± 0.15	76.52 ± 0.854	65.74 ± 0.897
	Medium	24.11 ± 0.103	2.54 ± 0.133	11.90 ± 0.071	2.03 ± 0.003	0.36 ± 0.005	2.36 ± 0.003	74.32 ± 0.140	74.76 ± 0.297	63.57 ± 0.413
	Coarse	23.72 ± 0.057	2.25 ± 0.035	10.51 ± 0.177	2.14 ± 0.003	0.27 ± 0.011	2.39 ± 0.003	72.28 ± 0.950	67.09 ± 0.420	57.51 ± 1.178
	Tender	25.82 ± 0.043	3.45 ± 0.015	12.60 ± 0.043	2.15 ± 0.005	0.42 ± 0.003	2.56 ± 0.005	75.47 ± 0.805	77.28 ± 0.279	66.14 ± 0.470
	Medium	24.38 ± 0.066	3.59 ± 0.075	12.31 ± 0.003	2.17 ± 0.003	0.57 ± 0.005	2.74 ± 0.003	74.51 ± 0.543	75.64 ± 0.337	63.99 ± 0.192
	Coarse	24.30 ± 0.034	3.24 ± 0.011	11.84 ± 0.035	2.16 ± 0.005	0.36 ± 0.005	2.52 ± 0.003	73.48 ± 0.287	69.02 ± 0.457	58.07 ± 0.337
SEM		0.091	0.038	0.084	0.002	0.004	0.003	0.766	0.379	0.583
CD 5%		NS	0.120	0.264	0.008	0.014	0.011	NS	NS	NS

Impact of gamma-rays on the growth performance and yield of S₁₃ mulberry variety

TABLE-3. Rearing efficiency of CB silkworm race (PMxNB₄D₂) fed with the leaves of S₁₃ mutant recovered at 6kR gamma irradiation (R₂ generation)

Treatments	Average larval weight (g)	Single cocoon weight (g)	Single shell weight (g)	Shell weight (%)	Filament length (m)	Reelability (%)	Renditta	Denier	Effective rate of rearing (%)
Control	3.33 ± 0.551	1.61 ± 0.029	0.25 ± 0.012	14.52 ± 0.092	663.64 ± 7.421	86.38 ± 1.085	8.05 ± 0.008	2.14 ± 0.088	78.08 ± 1.031
Mutant	3.56 ± 0.044	1.71 ± 0.027	0.28 ± 0.058	15.42 ± 0.244	722.73 ± 21.213	92.45 ± 2.205	7.45 ± 0.292	2.28 ± 0.035	86.68 ± 1.103
SEM	0.043	0.053	0.012	0.266	28.302	1.851	0.300	0.033	2.135
CD@5%	0.186	NS	NS	1.146	NS	7.966	NS	0.145	9.188