



## GENETIC VARIABILITY FOR DIFFERENT QUANTITATIVE TRAITS IN *DIANTHUS CARYOPHYLLUS* L. DURING MUTATION BREEDING

Rajib Roychowdhury and Jagatpati Tah

Cytogenetics & Plant Breeding Lab, Botany Department (UGC-CAS),  
The University of Burdwan, Burdwan-713104, West Bengal, India

### ABSTRACT

*Dianthus caryophyllus* L. is an important floricultural crop species throughout the world, specially in Eurasia. The present study was conducted to evaluate the nine *Dianthus* genotypes, including normal and mutagen treated, to assess the magnitude of variability and to pick up the heritable component of variation present in the quantitative traits. The analysis of variance revealed highly significant differences for all the characters studied, indicating the presence of substantial genetic variability. The field trial was followed Randomized Block Design with three replications for each genotypes. The phenotypic co-efficient of variation (PCV) was higher than genotypic co-efficient of variation (GCV) for all the characters studied, this shows the influence of the environmental effect on the characters. Highest GCV and PCV were noticed for total branches per plant and lowest for total no. of flowers per plant along with plant height (cm) at 50% flowering phase and flower diameter (cm) only in PCV. Broad sense heritability ranged from 33.33 (Days to seed germination) to 95.30 (Plant height at 50% flowering phase) per cent. Total leaves and branches showed high, plant height, days to branching and seeds per inflorescence showed moderate and flower per plant showed low genetic gain. Thus, these traits may serve as an effective selection parameter during breeding programme for crop improvement.

**KEY WORDS:** Carnation, Crop improvement, GCV, genetic gain, heritability, PCV.

### INTRODUCTION

*Dianthus caryophyllus*, commonly known as Carnation, is a popular cut-flower as per its floricultural impact throughout the world. A high agronomic demand related to yield production was noticed for this Caryophyllaceae member. Inducible mutation by chemical or physical agents can produce the development of *Dianthus* cultivars with more desirable floral characteristics and higher productivity (Roychowdhury, 2011; Roychowdhury and Tah, 2011). The knowledge of the extent to which the desirable characters are heritable is a prerequisite for any crop improvement programme, especially for mutation breeding. For this purpose, inducible mutation is a suitable source of producing variation through mutation breeding procedure (Domingo *et al.*, 2007) which can produce several improved mutant varieties with high demanding economic value (Din *et al.*, 2004). Various quantitative traits with agro-economical value like seed weight, number of branches, leaves, flowers, leaf area, etc., are very much complex in nature because they confirm polygenic inheritance and greatly influenced by minute fluctuation of environmental components. Genetic improvement of any crop is largely depending on the magnitude of several genetic parameters like analysis of variance of each mean value, phenotypic and genotypic variances, phenotypic and genotypic coefficients of variation (PCV and GCV), broad sense heritability and genetic gain on which the breeding methods are formulated for its further improvement. Analysis on genetic variability reveals about the presence of variation in their genetic constitution, and it is of utmost important as it provides the basis for effective selection. Wide spectrum of genetic variability has been induced in

*Dianthus caryophyllus* using both physical and chemical mutagens in order to utilize it in floricultural improvement and inheritance studies (Patil, 1966; Ashri, 1970; Gowda *et al.*, 1996). The extent of variability is measured by genotypic coefficient of variance (GCV) and phenotypic coefficient of variance (PCV) which provides information about relative amount of variation in different characters. Hence, to have a thorough comprehensive idea, it is necessary to have an analytical assessment of quantitative traits. Since heritability is also influenced by environmental factors, the information on heritability alone may not help in pin pointing characters enforcing selection. Nevertheless, the heritability estimates in conjunction with the predicted genetic gain will be more reliable (Johnson *et al.*, 1955). Heritability gives the information on the magnitude of inheritance of quantitative traits i.e. polygenic inheritance, while genetic gain will be helpful in formulating suitable selection procedures. Hence, the present study in *Dianthus* cultivar has been undertaken to study the available extent of genetic variation and the magnitude of heritability of several common quantitative characters and the maximum possible amount of genetic gain expected to occur during the selection programme for crop improvement.

### MATERIALS AND METHODS

The present study comprises of nine genotypes including normal (Control) and mutant of experimental plant material (*Dianthus caryophyllus* L.) after mutagenic treatment. Pure line seeds of this *Dianthus* variety were treated with three commonly used chemical mutagens namely Ethyl methane sulphonate (EMS), Sodium Azide (SA) and Colchicine (Col) with the concentrations of

0.1%, 0.4% and 0.7% for each mutagen (Roychowdhury and Tah, 2011a). The mutagen treated seeds were sown for raising first mutant ( $M_1$ ) generation and were maintained at Crop Research Farm (latitude 23.53° N, 22.56° S and longitude 83.25° E, 86° W), Botany Department, The University of Burdwan, West Bengal, India, during winter season (2009 to 2010). This sowing process followed Randomized Block Design (RBD) layout with three replications for each genotypes/treatments. Each genotype was sown in three rows of 5m length plot and 20 cm apart by adopting the spacing of 20 x 20 cm. Normal recommended cultural practices and plant protection measures were followed. The data were recorded on five randomly selected plants from each replication for quantitative traits studied like days to seed germination, plant height (cm) at 50% flowering phase, days to branching, total number of branches per plant, total leaves per plant, leaf area (cm<sup>2</sup>), total no. of flower per plant, diameter of flower (cm), seeds per inflorescence, weight (g) of 1000 seeds. The all data were analyzed statistically. For the analysis of genetic parameters, we consider the analysis of variance of each mean value, phenotypic and genotypic variances, phenotypic and genotypic coefficient of variation (PCV and GCV), broad sense heritability and genetic gain. Mean values were subjected to analysis of variance (ANOVA) to test the significance for each traits after Panse and Sukhatme (1957, 1967). Phenotypic and genotypic variances were estimated according to Lush (1940). The genotypic and phenotypic coefficients of variation (GCV and PCV) were worked out according to the method advocated by Singh and Chaudhary (1985).

Heritability in broad sense was worked out as per the methodology given by Allard (1960). Expected genetic gain (GG) of the genotypes and its per cent of mean at 5% intensity of selection pressure (2.06 after Kang *et al.*, 1983) were calculated according to Singh and Chaudhary (1985).

## RESULTS AND DISCUSSION

In this investigation, all 10 studied quantitative characters showed significant differences in the mean sum of squares due to genotypes or treatments (Table 1) which indicate that the nine genotypes of *Dianthus* cultivar including the control and mutagen treated ones were genetically divergent. So, there is a huge scope for selection of promising mutant lines with different metrical traits from the present gene pool. The presence of wide range of variability might be due to diverse source of materials after mutation taken as well as environmental influence affecting the phenotypes. The calculated value of variance ratio was significant at 5 % level in case of days to seed germination, leaf area and weight of 1000 seeds; other characters showed significant at 1 % level of significance. The significant critical difference (CD) value indicates that *Dianthus* cultivar was no doubt suitable in the location where prevailing agro-climatic factors provided plantation of crop was in proper time. Higher the CD value indicates higher the stability in that climate mutagen (Roychowdhury and Tah, 2011a). Here, total leaves per plant shows higher CD value (4.49), days to branching and seeds/inflorescence show moderate (2.74 and 2.64, respectively) and weight of 1000 seeds represents lower CD value (0.21).

**TABLE 1.** Analysis of variance for 10 quantitative characters in 9 genotypes of *Dianthus caryophyllus* including normal control and mutant ones grown at Crop Research Farm, Burdwan University, during winter (2009-2010). Mean sum square (MSS) of replications, treatments and error along with critical difference (CD) and coefficient of variation (CV) were calculated for each studied characters. 'df' denotes degree of freedom. '\*' and '\*\*' sign indicate the significant at probability 5% and 1% level of significance, respectively.

| Source                        | df | Quantitative characters studied |                                          |                   |                             |                    |                              |                           |                         |                     |                          |
|-------------------------------|----|---------------------------------|------------------------------------------|-------------------|-----------------------------|--------------------|------------------------------|---------------------------|-------------------------|---------------------|--------------------------|
|                               |    | Days to seed germination        | Plant height (cm) at 50% flowering phase | Days to branching | Total no. of branches/plant | Total leaves/plant | Leaf area (cm <sup>2</sup> ) | Total no. of flower/plant | Diameter of flower (cm) | Seeds/inflorescence | Weight (g) of 1000 seeds |
| Replication                   | 2  | 0.15                            | 0.286                                    | 1.148             | 1.78                        | 0.704              | 0.04                         | 3.37                      | 0.02                    | 1.037               | 0.001                    |
| Treatment                     | 8  | 0.37*                           | 32.698**                                 | 13.759**          | 3.171**                     | 105.12**           | 0.69*                        | 10.62**                   | 0.106**                 | 70.79**             | 0.026*                   |
| Error                         | 16 | 0.148                           | 0.528                                    | 1.315             | 0.359                       | 3.537              | 0.258                        | 1.287                     | 0.03                    | 3.62                | 0.015                    |
| Coefficient of Variation (CV) |    | 8.926                           | 1.437                                    | 6.686             | 12.535                      | 3.132              | 13.262                       | 3.119                     | 4.214                   | 10.01               | 7.953                    |
| Critical Difference (CD)      |    | 0.66                            | 1.73                                     | 2.74              | 1.43                        | 4.49               | 0.88                         | 1.96                      | 0.41                    | 2.64                | 0.21                     |

The estimation on genetic parameters like genotypic ( $\sigma^2_g$ ) and phenotypic ( $\sigma^2_p$ ) variance, genotypic (GCV) and phenotypic (PCV) coefficient of variation, broad sense heritability ( $h^2_{bs}$ ) and genetic gain (GG) of the population of *Dianthus* genotype under study is given in Table 2. A wide range of variation was observed among different genotypes/treatment with regard to different characters. The range of mean within the character was generally in equilibrium for most characters except for total leaves per

plant, total number of flower per plant and no. of seeds per inflorescence.

The maximum genotypic and phenotypic variation were obtained for total leaves per plant and plant height (cm) at 50% flowering phase, while moderate variation was observed for days to branching, seeds per inflorescence and total no. of flower per plant. This indicated that the environment did not influence these characters very much. Values of phenotypic and genotypic variance were very

much close for weight (g) of 1000 seeds and total no. of branches per plant. The character with almost equal value of phenotypic and genotypic variance can be considered stable. Low level of genotypic and phenotypic variance for days to seed germination, leaf area (cm<sup>2</sup>), flower diameter

(cm) which also shows negative value for genotypic variance and 1000 seed weight (g) is indicative of stable nature of these characters. Similar findings were reported by Ganesan *et al.* (1994) and Rao *et al.* (1996).

**TABLE 2.** Component of genetic parameters for 10 quantitative characters among nine *Dianthus caryophyllus* genotypes.

| Sl No. | Quantitative Traits                      | Mean (X) | Range |       | Components of Variance      |                             |                             | GCV   | PCV   | h <sup>2</sup> <sub>bs</sub> % | GG     | GG as % of mean |
|--------|------------------------------------------|----------|-------|-------|-----------------------------|-----------------------------|-----------------------------|-------|-------|--------------------------------|--------|-----------------|
|        |                                          |          | Max   | Min   | σ <sup>2</sup> <sub>g</sub> | σ <sup>2</sup> <sub>p</sub> | σ <sup>2</sup> <sub>e</sub> |       |       |                                |        |                 |
| 1.     | Days to seed germination                 | 4.31     | 5.01  | 4.13  | 0.074                       | 0.222                       | 0.148                       | 0.235 | 0.406 | 33.33                          | 0.185  | 4.29            |
| 2.     | Plant height (cm) at 50% flowering phase | 50.57    | 55.1  | 45.6  | 10.723                      | 11.251                      | 0.528                       | 0.239 | 0.246 | 95.30                          | 6.43   | 12.72           |
| 3.     | Days to branching                        | 17.15    | 22.6  | 13.21 | 4.148                       | 5.463                       | 1.315                       | 0.439 | 0.505 | 75.93                          | 3.186  | 18.58           |
| 4.     | Total no. of branches/plant              | 4.78     | 7.4   | 3.51  | 0.937                       | 1.296                       | 0.359                       | 0.750 | 0.882 | 72.30                          | 1.441  | 30.15           |
| 5.     | Total leaves/plant                       | 60.04    | 85.91 | 34.34 | 33.861                      | 37.398                      | 3.537                       | 0.359 | 0.377 | 90.54                          | 10.852 | 18.07           |
| 6.     | Leaf area (cm <sup>2</sup> )             | 3.83     | 4.86  | 2.82  | 0.144                       | 0.402                       | 0.258                       | 0.367 | 0.613 | 35.82                          | 0.28   | 7.31            |
| 7.     | Total no. of flower/plant                | 36.37    | 43    | 25.4  | 3.111                       | 4.398                       | 1.287                       | 0.179 | 0.214 | 70.74                          | 2.57   | 7.06            |
| 8.     | Diameter of flower (cm)                  | 4.11     | 4.6   | 3.5   | -0.07                       | 0.029                       | 0.099                       | ---   | 0.154 | 241.37                         | ---    | ---             |
| 9.     | Seeds/inflorescence                      | 19       | 27.6  | 12.32 | 3.972                       | 5.562                       | 1.59                        | 0.561 | 0.638 | 71.41                          | 2.932  | 15.43           |
| 10.    | Weight (g) of 1000 seeds                 | 1.54     | 1.82  | 1.25  | 0.011                       | 0.026                       | 0.0147                      | 0.253 | 0.388 | 42.30                          | 0.092  | 5.97            |

#### Legends:

σ<sup>2</sup><sub>g</sub> = genotypic variance, σ<sup>2</sup><sub>p</sub> = phenotypic variance, σ<sup>2</sup><sub>e</sub> = environmental variance, GCV = genotypic coefficient of variance, PCV = phenotypic coefficient of variance, h<sup>2</sup><sub>bs</sub> = broad sense heritability, GG = genetic gain.

In general, phenotypic coefficient of variation (PCV) was higher in magnitude than the genotypic coefficient of variation (GCV) for all the characters studied, the close resemblance between the corresponding estimates of both PCV and GCV in almost all the characters suggested that the environment had little effect in those characters expression (Jalgaonkar *et al.*, 1990). The GCV provides a measure to compare genetic variability present in various quantitative characters. The highest value of it was recorded for total branches per plant, moderate for days to branching and seeds per inflorescence and lowest for total flowers per plant. Flower diameter (cm) showed no GCV because genotypic variance was negative. The higher value clearly indicated high degree of genotypic variability in these quantitative traits in *Dianthus caryophyllus*. PCV which measure total relative variation was highest for total branches per plant, moderate for seeds per inflorescence, leaf area, days to branching and seed germination, lowest for flower diameter (cm), total no. of flowers per plant and plant height (cm) at 50% flowering phase. Similar result was reported for plant height (cm) Pathania *et al.* (1988). High values of GCV suggest better improvement scope for these traits by selection. However, the estimation of heritable variation with the help of genetic coefficient of variation alone may be misleading. Burton (1951, 1952) suggested that the genetic coefficient of variation together with heritability

estimates gave the better picture of the extent of heritable variation.

Heritability (h<sup>2</sup>) and genetic gain (GG) estimates were interpreted as low, medium and high as per the classification of Johnson *et al.* (1955). Broad sense heritability (h<sup>2</sup><sub>bs</sub>) ranged from 33.33 (Days to seed germination) to 95.30 (Plant height at 50% flowering phase) per cent. High heritability was recorded for plant height at 50% flowering phase (95.3%) and total leaves per plant (90.54%), moderate heritability for days to branching (75.93%), total branches per plant (72.3%), seeds per inflorescence (71.41%) and total flowers per plant (70.74%), lowest for days to seed germination (33.33%) and leaf area (35.82%). Flower diameter showed negative value due to its genotypic variance. High heritability combined with high genetic gain as per cent of mean was observed for total leaves per plant and total no. of branches per plant. This indicates the lesser influence of environment in expression of these characters and prevalence of additive gene action in their inheritance, hence amenable for simple selection. High heritability with moderate genetic gain as per cent of mean was recorded for plant height at 50% flowering, days to branching and seeds per inflorescence indicating that these characters were governed by additive gene interaction. High heritability coupled with low genetic gain as per cent of mean was recorded for total flowers per plant indicating non-additive gene action for these traits.

**CONCLUSION**

The knowledge on heritability of traits is helpful to decide the selection procedure to be followed to improve the trait in a situation. Higher estimates of heritability with higher genetic gain as per cent of mean was observed for total leaves per plant and total no. of branches per plant indicating the presence of additive gene action and so selection can be easily done for these traits. The trait which expressed high heritability and low genetic gain showed non additive gene interaction, hence heterosis breeding would be recommended for that trait.

**REFERENCES**

- Allard, R.W. (1960) Principles of plant breeding. pp. 89-98. John Wiley and Sons, New York.
- Ariyo, O.J. (1987) Variation and Heritability of Fifteen Characters in Okra (*Abelmoschus esculentus* L.). *Tropical Agriculture* **34**, 67 – 71.
- Ashri, A. (1970) A dominant mutation with variable penetrance and expressivity induced by diethyl sulfate in peanuts, *Arachis hypogaea* L. *Mutation Res.* **9**, 473-480.
- Burton, G.W. (1951) Quantitative inheritance in Pearl millet (*P. glaucum*). *Agron. J.* **43**, 409-417.
- Burton, G.W. (1952) Quantitative inheritance in grasses. *6th Int. Grassland Congress* **1**, 277-83.
- Din, R., Qasim, M. and Ahmad, K. (2004) Radio sensitivity of various wheat genotypes in M<sub>1</sub> generation. *Int. J. Agri. Biol.* **6**, 898-900.
- Domingo, C., Andres, F. and Talon, M. (2007) Rice cv. Bahia mutagenized population: a new resource for rice breeding in the Mediterranean basin. *Span. J. Agric. Res.* **5**, 341-347.
- Galeta, N., Mohammed, H. and Zelleke, H. (2005) Genetic variability and genetic advance in sorghum (*Sorghum bicolor* (L.) Moench) germplasm. *Crop Res.* **30**, 439-445.
- Gowda, M.V.C., Nadaf, H.L. and Sheshagiri, R. (1996) The role of mutation in intraspecific differentiation of groundnut (*Arachis hypogaea* L.). *Euphytica* **90**, 105-113.
- Ganesan, K. (1994) Genetic studies in F<sub>2</sub> and F<sub>3</sub> of tall x dwarf rice crosses. *Madras Agricultural Journal* **81**, 30-32.
- Jabeen, N., Ahmed, N. and Tanki, M.I. (1998) Genetic variability in hot pepper (*Capsicum annum* L.). *Agric. Sci. Digest* **18**: 23-26.
- Jalgaonkar, R., Jamadagni, B.M. and Salvi, M.J. (1990) Genetic variability and correlation studies in turmeric. *Indian Cocoa, Arecanut and spices Journal* **15** (1), 20-22.
- Johnson, H.W., Robinson, H.F. and Comstock, R.E. (1955) Estimation of genetic and environmental variability in soybean. *Agronomy Journal* **47**, 314-318.
- Kang, M.S., Mille, J.D. and Tai, P.Y.P. (1983) Genetic and phenotypic path analysis and heritability in sugarcane. *Crop Sci.* **23**, 643-647.
- Lush, J.L. (1940) Intra - sire correlation and regression of offspring on dams as a method of estimating heritability of characters. *Proc. Amer. Soc. Animal Production* **33**, 293-301.
- Munshi, A.D. and Behera, T.K. (2000) Genetic variability, heritability and genetic advance for some traits in chillies (*Capsicum annum* L.). *Vegetable Sci.* **27**, 39 – 41.
- Nayak, A.R., Chaudhary, D. and Reddy, J.N. (2002) Genetic variability, heritability and genetic advance in scented rice. *Indian Agriculturist* **46**, 45-47.
- Panse, V.G. (1957) Genetics of quantitative characters in relation to Plant Breeding. *Ind. J. Genet. & Plt. Breed.* **17**, 318-28.
- Panse, V.G. and Sukhatme, P.V. (1967) Statistical Methods for Agricultural Workers. pp. 381. ICAR publication, New Delhi. 2<sup>nd</sup> Edition.
- Pathania, N., Singh, M. and Arya, P.S. (1988) Genetic evaluation of some economic traits in turmeric (*Curcuma longa* L.). *Himachal Journal of Agricultural Research* **14** (1), 38-43.
- Patil, S.H. (1966) Mutations induced in groundnut by X-rays. *Indian J. Genet.* **26A**, 334-348.
- Rao, T.P., Gomathinayagam, P. and Soundrapandian, S. (1996) Genetic variability and character association studies in semi-dry rice. *Madras Agricultural Journal* **83**(3), 185-188.
- Roychowdhury, R. (2011) Effect of Chemical Mutagens on Carnation (*Dianthus caryophyllus* L.): A Mutation Breeding Approach, LAP Lambert Academic Publishing, Germany.
- Roychowdhury, R. and Tah, J. (2011) Mutation breeding in *Dianthus caryophyllus* for economic traits. *Elec. J. Plt. Breed.* **2**(2), 282-286.
- Roychowdhury, R. and Tah, J. (2011a) Chemical mutagenic action on seed germination and related agro-metrical traits in M<sub>1</sub> *Dianthus* generation. *Current Botany* **2**(8), 19-23.
- Singh, R.K. and Chaudhary, B.D. (1985) Biometrical Methods in Quantitative Genetic Analysis. pp. 318. Kalyani Pub., Ludhiana.