



EFFECTS OF CRUDE OIL AS A SOIL CONTAMINANT ON SEEDLING GROWTH OF *JATROPHA CURCAS* L.

¹Agbogidi, O.M., ²Ilondu, E.M. and ¹Ohwo, O.A.

¹Department of Forestry and Wildlife, Faculty of Agriculture,

²Department of Botany, Delta State University, Asaba Campus, Nigeria.

ABSTRACT

A study was conducted in 2010 to evaluate the effects of crude oil as a soil contaminant on the performance of *Jatropha curcas* seedlings in Asaba, Delta State, Nigeria. Oil concentrations used were 0.0, 2.0, 4.0, 6.0, 8.0 and 10.0% w/w. The experiment was arranged in a randomized complete block design with four replicates and monitored for 10 weeks after transplanting. The results showed that contamination of soil with crude oil significantly reduced ($P \geq 0.05$) the performance of *Jatropha curcas* seedlings as regards plant height, number of leaves, leaf area and collar diameter when compared with seedlings grown in the uncontaminated soils. The performance of the seedlings was oil concentration dependent. Other symptoms observed at the higher levels of oil treatments were yellowness of leaves, leaf fall/drop, chlorosis, growth retardation and stagnation as well as wilting. This study has shown that crude oil in soil has a highly significant effect of reducing the performance of *J. curcas* seedlings. The oil level most tolerated by the plant is 2.0% w/w and above this level, significant reductions were recorded. This study has great implications on the sustainability of this multipurpose species, which has been identified as a potential biofuel species.

KEYWORDS: performance, *Jatropha curcas* seedlings, crude oil, soil.

INTRODUCTION

Jatropha curcas is a perennial shrub that is native to Central America. It is now widely cultivated in tropical and subtropical regions of the world becoming even naturalized in other areas (Achtein, 2008). Commonly called Barbados nut, purging nut or physic nut, the plant produces flowers and fruits throughout the year. It is resistant to a high degree of acidity (Henry, 2009). The fruit contains 37.5% shell and 62.5% seeds (Jepsen *et al.*, 2006). Seeds resemble castor in shape, black in colour and are 42% husk and 58% kernel. The seeds contain 27-40% oil that can be processed to produce a high quality biodiesel fuel useable in a standard diesel engine (Biswas *et al.*, 2006; Kureel, 2006; Wini *et al.*, 2006). Achten *et al.* (2008) noted that *J. curcas* is mainly cultivated for extraction of biodiesel and is one of the best sources of biodiesel. Achten *et al.* (2008) further maintained that one hectare of *J. curcas* yields 6-7MT of seeds while one tonne of *J. curcas* seeds yields 30kg oil product and 700kg oil cake. The rapid growth and development of *J. curcas* will represent the most immediate and available response of coping with depleting oil reserves, meeting growing energy demand in developing countries like Nigeria *i.e.* to support development in rural areas. The renewed interest in *Jatropha* cultivation could also reduce dependence on a limited number of exporting nations by diverting their energy sources and supplies as well as to challenge emerging economies in tropical regions to supply the global energy market with competitive price liquid biofuel besides stimulating economic growth, improving the environment and reducing dependency on imported oil (Kureel, 2006; Tigere *et al.*, 2006). Economically, *J. curcas* is a medicinal plant: the seeds prevent constipation, the sap for wound healing and bleeding, the leaves as tea

against malaria, and a potential source of herbal drugs in dental complaints. The plant also has the possibility of reclaiming marginal soils by reanchoring the soil with its substantial root and reduces possibility of erosion (Jepsen *et al.*, 2006; Openshaw, 2000). Openshaw (2000) also reported that *J. curcas* is planted in form of hedges around gardens or fields to protect the crops against roaming plants to demarcate boundaries of fields and homesteads and as a source of shade for coffee plants in India and Cuba (Kumar *et al.*, 2008a). Although *J. curcas* grows best in sandy well – drained soils, it also does well in warmer areas, as it can withstand severe heat, light frost for not too long, very poor soils and grows in saline conditions (Kumar *et al.*, 2008b). Food production and transport are particularly dependent on diesel engines and biodiesel is a crucial part of bringing energy security benefits to the global food supply. In times of international terrorism there are heightened concerns about nuclear proliferation and energy security argues that the future must belong to bioenergy. If we can achieve energy security, we can not only free ourselves from dependence on fossil fuels, we can also realize environmental security and a whole host of other central developmental and poverty alleviation goals. This is because decreasing oil production from almost all the oil reserves is attracting the world's eyes towards renewable energy sources. By promoting a shift towards low carbon economic activities, governments can not only help avoid dangerous climate change impacts but can also make the region more competitive, contributing to a faster recovery from the current economic slowdown. The oil industry needs to be prepared for tomorrow, even in these uncertain times *Jatropha* and other non-feed crops produce biofuels and hold particular promise for sustainable development and a

sustainable environment (Forson *et al.*, 2004). Its use is considerably less expensive feedstock, gives the producer a significant cost advantage over oil petroleum products. *Jatropha* and other non-food oil-bearing crops are the cheapest and the most viable business propositions. *Jatropha* can bring significant environmental benefits. It can replace jet fuel and diesel from petroleum without interfering with food crops or leading to the clearing of forests (Sukarin *et al.*, 1987). The good thing about the plant is that a tree shrub that lives for a longtime, doing its job, producing oil and sequestering lots of CO₂ from the atmosphere is produced. It can as well help to increase income from plantations and agro-industries besides helping to increase rural incomes, self-sustainability and alleviation of poverty. Biodiesel produced from *Jatropha* is one of the most promising solutions for tackling the growing carbon emissions from transport. *Jatropha* cultivation has a huge potential for replication worldwide improving the livelihood of many more (Biswas *et al.*, 2006). Presently, *J. curcas* found in Nigeria are those traditionally used as hedging plants and grave markers. Most recently, research institutes and investors have started establishing plantation of the plant because of its potential biodiesel production (Onyeme *et al.*, 2008; Ige *et al.*, 2010). The ability of *J. curcas* to grow in gravelly, sandy, degraded, acidic, poor/marginal soil according to Openshaw (2000) and Jepsen *et al.* (2006) has provided a basis for this investigation.

Crude oil pollution is an inevitable phenomenon in oil producing and consuming areas worldwide (Agbogidi and Eshegbeyi, 2006). It stems from human error, accidental discharge and other sources (Agbogidi and Ayelo, 2010). The severity of oil pollution effects varies with the quantity, plant species, age of the plant, adequacy of the responses as well as other factors (Vwioko *et al.*, 2006; Agbogidi, 2009a). Oil pollution has deleterious effects on soil hence significant effects on plant growth. Although literatures abound on the susceptibility of plant species to crude oil and its products (Anoliefo and Edegbai, 2000; Agbogidi *et al.*, 2006; Anoliefo *et al.*, 2006; Bamidele *et al.*, 2007; Agbogidi, 2009b), there is scarcity of documented information on the influence of crude oil on *J. curcas*. It is against this background that a study as this has been embarked upon to provide information on the performance of *J. curcas* seedlings as affected by crude oil in soil with a view to establishing the critical level this valuable multipurpose species can be significantly affected.

MATERIALS AND METHODS

The study was conducted in 2010 at latitude 6°14'N and longitude 6°49'E (Asaba Meteorological Office, 2010) at the nursery site of the Department of Forestry and Wildlife, Delta State University, Asaba Campus, Nigeria. Matured fruits were harvested from the parent plants in Asaba, Delta State. The fruits were depulped mechanically to extract the seeds. Healthy seeds were selected and sorted out. Viability tests were carried out on the seeds using floatation technique. The depulped seeds (600) in number were sown in the Departmental nursery and the basic nursery techniques were observed. The soil treatment was thoroughly mixed with appropriate crude oil levels

before the polypots (10/15cm in dimension) were each filled up with the oil-soil mixture. Oil concentrations used were 0.0, 2.0, 4.0, 6.0, 8.0 and 10.0% w/w. The soil sample was obtained as a pooled sample from the Gmelina plantation behind the Departmental office. The soil was air-dried and passed through a 2mm sieve. The crude oil was obtained from the Nigeria National Petroleum Corporation (NNPC), Warri., Delta State. The seedlings (6 weeks of age) were transplanted into the oil-contaminated soil and the uncontaminated soil (control) in the polypots and watered immediately to field capacity and after wards, every other day till the end of the experiment. There were therefore six treatments, replicated four times and arranged in randomized complete block design. One seedling was transplanted into each polypot and each treatment comprised 10 polypots indicating a total of 180 seedlings of relatively the same height were transplanted. The set-up was monitored for 10 weeks after transplanting (WAT) while growth parameters were measured fortnightly with effect from the second week after transplanting. Plant height was determined with a meter rule at the distance from soil level to the terminal bud; number of leaves was by visual counting of the leaves, leaf area was measured by tracing the margins of the leaf on a graph paper and the total leaf area/plant was obtained by counting the leaf number of 1-cm squares following the procedure of Agbogidi and Eshegbeyi (2006). The collar diameter was determined by measuring at two centimeter above soil level with vernier calipers. Data obtained were exposed to one-way analysis of variance (ANOVA) and the significant means were separated with the Duncan's Multiple Range Tests (DMRT) using SAS (2005).

RESULTS AND DISCUSSION

As shown in Tables 1, 2, 3 and 4, contamination of soil with crude oil significantly ($P \geq 0.05$) reduced all the growth indices assessed. Plant height, number of leaves, leaf area and collar diameter were significantly affected by oil treatment at ($P \geq 0.05$) when compared with seedlings grown in the uncontaminated soils. The growth of seedlings was progressively poorer as the level of oil increased indicating that the performance of the seedlings was oil concentration-dependent. For example, while 25.8cm and 22.9cm height values were recorded for seedlings grown in 0.0 and 2.0% w/w of oil at 6 WAT, 16.9cm and 16.6cm were obtained for seedlings exposed to 8.0 and 10.0% w/w of the oil concentration respectively. The same trend was also observed in number of leaves, leaf area and collar diameter (Tables 2, 3 and 4) respectively. Other visible symptoms were yellowness of leaves, leaf drop, chlorosis, necrosis growths retardation and stagnation as well as wilting at the highest level (10.0% w/w) of crude oil treatment of soil. These observations are similar to the reports of Achuba (2006), Anoliefo *et al.* (2006), Agbogidi and Dolor (2007), Agbogidi *et al.* (2007), Agbogidi (2009b), Agbogidi (2010) on groundnut, cowpea, common beans, dikanut, maize, Gmelina and soyabeans respectively. Yellowness of leaves could be due to nutrient immobilization as oil pollution has been reported to cause unavailability of some essential nutrients while some toxic ones may be more readily available (De Jong, 1980; Benka-Coker and

Ekundayo, 1995; Benka–Coker and Ekundayo, 1997). The observed leaf drop/fall and wilting could be linked to the inability of the seedlings to absorb water because they were watered like their counterparts in the uncontaminated subplots. This could have stemmed from the alteration caused by the presence of the oil in soil and this observation is in agreement with the findings of Atuanya (1987) and Ekundayo and Obuekwe (1997) that oil in soil affects the physical, chemical and biological properties of the soil. Growth retardation and stunting could have been attributed to both soil-plant-water interrelations as well as a disruption in the xylem and phloem vessels due to the adulterated soil structure and this could have culminated in the over growth stagnation in seedlings grown in the highest level of oil contamination in the highest level of oil

contamination. This observation lays credence to the reports of Agbogidi and Ofuoku (2005) and Agbogidi and Eshegbeyi (2006) that oil level seemed to exert the greatest influence on plant growth and yields. The presence of other toxic components of the crude oil as well as heavy metals could have also accounted for the observed growth reduction, stunting and stagnation in the highest levels of oil application to soil as observed in the present study. This study has shown that crude oil in soil has a highly significant effect of reducing the performance of *J. curcas* seedling in terms of plant height, number of leaves, leaf area and collar diameter. The oil level most tolerated (critical) by the plant is 2.0% w/w above this level, significant reductions were recorded.

TABLE 1. Plant height (cm) of *Jatropha curcas* seedlings grown in soil contaminated with crude oil

Oil in soil	Plant height/WAT					
(% w/w)	2	4	6	8	10	Means
0.0	18.9	20.4	23.6	25.8	26.7	23.1a
2.0	18.6	20.2	22.9	25.3	25.8	22.6b
4.0	18.0	18.3	18.4	18.6	18.8	18.4c
6.0	17.4	17.5	17.6	17.7	17.8	17.6d
8.0	17.1	17.2	17.3	16.9	16.8	17.6d
10.0	16.9	17.0	16.8	16.6	16.5	16.8e
Means	17.8	18.4	19.4	20.2	20.0	

Means with different letters are significantly different at $P \leq 0.05$ level by Duncan's Multiple Range Tests.

TABLE 2. Number of leaves of *Jatropha curcas* seedlings as affected by crude oil contaminated of soil

Oil in soil	Number of leaves/WAT					
(% w/w)	2	4	6	8	10	Means
0.0	5.1	6.3	7.8	8.5	9.7	7.3a
2.0	5.0	6.2	7.6	8.4	9.6	7.4a
4.0	4.8	5.2	5.4	5.6	5.7	5.3b
6.0	4.7	5.0	5.1	5.1	5.1	5.0c
8.0	4.6	4.3	4.2	4.1	4.0	4.2d
10.0	4.3	4.2	4.1	4.0	3.8	4.1e
Means	4.75	5.20	5.70	5.95	6.32	

Means with different letters are significantly different at $P \leq 0.05$ level by Duncan's Multiple Range Tests.

TABLE 3. Leaf area (cm^2) of *Jatropha curcas* seedlings as influence by crude oil in soil

Oil in soil	Leaf area/WAT					
(% w/w)	2	4	6	8	10	Means
0.0	126.4	267.2	324.4	386.6	524.3	325.8a
2.0	120.2	254.1	320.2	382.2	519.1	319.2b
4.0	118.2	225.6	260.2	286.3	271.4	232.3c
6.0	115.1	201.7	214.6	216.7	219.2	193.5c
8.0	110.0	126.9	114.7	112.6	100.2	112.9e
10.0	101.0	118.1	100.1	95.5	90.3	101.0f
Means	115.2	198.9	222.4	214.5	287.4	

Means with different letters are significantly different at $P \leq 0.05$ level by Duncan's Multiple Range Tests.

TABLE 4. Collar diameter (cm) of *Jatropha curcas* as affected by crude oil contamination of soil

Oil in soil (% w/w)	Collar diameter /WAT					Means
	2	4	6	8	10	
0.0	0.73	0.90	1.10	1.31	1.49	1.11a
2.0	0.73	0.86	1.09	1.20	1.38	1.05a
4.0	0.71	0.74	0.82	0.89	1.04	0.84b
6.0	0.70	0.73	0.80	0.82	1.00	0.81c
8.0	0.70	0.66	0.64	0.60	0.58	0.63d
10.0	0.69	0.64	0.62	0.54	0.52	0.60e
Means	0.71	0.76	0.85	0.90	1.00	

Means with different letters are significantly different at $P \leq 0.05$ level by Duncan's Multiple Range Tests.

REFERENCES

- Achten, W.W., Verchot, L., Franken, Y.J., Mathijs, E., Sighn, V.P., Aert, R. and Muys, B. (2008) *Jatropha* biodiesel production. *Bioenergy* 32 (12): 1063-1084.
- Agbogidi, O.M. and Eshegbeyi, O.F. (2006) Performance of *Dacryodes edulis* (Don. G. Lam H.J.) seeds and seedlings in a crude oil contaminated soil. *Journal of Sustainable Forestry* 22 (3/4): 1-14.
- Agbogidi, O.M. (2009a) Yield components of six cultivars of cowpea (*Vigna unguiculata* (L.) Walp) grown on soil contaminated with spent engine oil. *Acta Agronomica Nigeriana*, 9 (1 and 2): 1-6.
- Agbogidi, O. M. (2009b) Effects of soil contaminated soil with spent lubricating oil on the germination of *Gmelina arborea* (Roxb.) seeds. *African Journal of Natural Sciences* 2 (4): 11-14.
- Agbogidi, O.M. (2009c). Locational effect on the performance of maize (*Zea mays* L) in soils treated with crude oil. *African Journal of Environmental Pollution and Health* 7(1): 36-42.
- Agbogidi, O.M. (2010). Screening six cultivars of cowpea (*Vigna unguiculata* (L) Walp) for adaptation to soil contaminated with spent engine oil. *Academic Arena* 2(4): 33-40.
- Agbogidi, O.M. and Ayelo, E. (2010) Germination of African oil bean (*Pentaclethra macrophylla*, Benth.) Seeds grown in crude oil polluted soil. In: Onyekwelu, J., Adekunle, V.A.J. and Oke, D.O. (eds.). Proceedings of the 2nd biennial national conference of the forests and forest products society held at the Federal University of Technology, Akure, Nigeria between 26th and 29th of April 2010. Pp 105-111..
- Agbogidi, O.M. and Dolor, D. E. (2007) An assessment of the growth of *Irvingia gabonensis* (Aubry-Lecomte Ex O' Rorte) Bail seedlings as influenced by crude oil contamination of soil. *Asian Journal of Plant Sciences* 2: 1287-1292.
- Agbogidi, O.M., Eruotor, P.G. and Akparaobi, S.O. (2007) Effects of crude oil levels on the growth of maize (*Zea mays* L). *American Journal of Food Technology* 2(6): 529-535.
- Agbogidi, O. M. and Ofuoku, A.U. (2005) Response of sour sop (*Annona muricata* Linn.) to crude oil levels. *Journal of Sustainable Tropical Agricultural Research* 16: 98-102.
- Anoliefo, G. O. and Edegbai, B. O. (2000) Effect of spent engine oil as a soil contaminant on the growth of two egg plant species, *Solanum melongena* L. and *S. incanum* L. *Journal of Agriculture, Forestry and Fisheries* 1:21-25.
- Anoliefo, G. O. and Vwioko, D. E. (2001) Tolerance of *Chromolaena odorata* (L) K. and R. grown in soil contaminated with spent lubricating oil. *Journal of Tropical Bioscience* 1(1): 20-24.
- Anoliefo, G. O., Isikhuemhen, O. and Ohimain, E. (2006) Sensitivity studies of the common bean (*Vigna unguiculata*) and maize (*Zea mays*) to different soil types from the crude oil drilling site at Kutchalli, Nigeria. *Journal of Soils and Sediments* 6 (1): 30-36.
- Aregheore, E.M., Becker, K. and Mallar, H.P. (2003) Detoxification of a toxic variety of *Jatropha curcas*. *South Pacific Journal of Natural Science* 21:50-56.
- Atuanya, E.I. (1987). Effect of waste engine oil pollution on physical and chemical properties of soil: a case study of waste contaminated Delta soil in Bendel State. *Nigerian Journal of Applied Sciences* 5: 155-176.
- Asaba Meteorological Station (2010) National Meteorological Bulletin 2010 Lagos, Nigeria.
- Awe, A.B. Baiyewu, R.A., Fatade, B.O., Adeleke, T.O., Omidran, M.B and Salaudeen, G.T. (2010) Photochemical and antimicrobial evaluation of the leaves of *Jatropha curcas*. In: Onyekwelu, J., Adekunle, V.A.J. and Oke, D.O. (eds.). Proceedings of the 2nd biennial national conference of the forests and forest products society held at the Federal University of Technology, Akure, Nigeria between 26th and 29th of April 2010. Pp 445 – 449.
- Bamidele, J. F., Agbogidi, O.M., and Ohienbor, O.M. (2007) Phyto-toxic effects of gasoline on *Ischaemum rugosum* (Salisb): a wetland species. *American Journal of Plant Physiology* 2(4): 276-281.
- Benka-Coker, M. O. and Ekundayo, J. A. (1995) Effect of an oil spill on soil physicochemical properties of a spill site in the Niger Delta area of Nigeria. *Environmental Monitoring and Assessment* 36:93-104.

- Benka – Coker, M. O. and Ekundayo, J. A. (1997) Applicability of evaluating the ability of microbes isolated from an oil spill site to degrade oil. *Environmental Monitoring and Assessment* 45: 259 – 272.
- Biswas, S., Kaushik, N. and Srikanth, G. (2006) Biodiesel: technology and business opportunities an insight presented at the Biodiesel conference Towards Energy Independence – Focus on *Jatropha*. Rashtrapati Nilayam, Bolaram, India.
- De Jong, E. (1980) The effect of a crude oil spill on cereals. *Environmental Pollution* 22: 187-196.
- Ekundayo, E. O. and Obuekwe, C. A. (1997) Effects of oil spill on soil physico-chemical properties of a spill site in a typical paleodult of Midwestern Nigeria. *Environmental Monitoring Assessment* 45:209-221.
- Forson, F.K., Oduro, E.K. and hammon Donkoh, E. (2004) Performance of *Jatropha* oil blends in a diesel engine. *Renewable Energy* 29:1135-1145.
- Henry, R. (2009) Plant resources for food, fuel and conservation. Earthsacn.
- Jepsen, J.K., Henning, R.K. and Nyati, B. (2006) Generative Propagation of *Jatropha curcas* L. on Kalahari sand. *Environment Africa, Zimbabwe*.
- Kumar, A., Ashwani, C.A. and Satyawati, S. (2008a) An evaluation of multipurpose oil seed crop for industrial uses (*Jatropha curcas* L.): a review. *Industrial Crops and Products* 1-8.
- Kumar, G.P., Yadav, S.K., Thawale, P.R., Singh, S.K. and Juwarka, A. A. (2008b) Growth of *Jatropha curcas* on heavy metal contaminated soil amended with industrial wastes and Azotobacter: a greenhouse study. *Bioresource Technology* 99:2078-2082.
- Kureel, R.S. (2006) Prospects and potentials of *Jatropha curcas* for biodiesel production. A paper presented at the Biodiesel conference Towards Energy Independence – Focus on *Jatropha*. Rashtrapati Nilayam, Bolaram, India.
- Makkar, H.P., Becker, K., Sporer, F. and Wink, M. (1997). Studies on nutritive potentials n toxic constituents of fiddler provenances of *Jatropha curcas*. *Journal of Agricultural of Food Chemistry* 45:3152-3157.
- Openshaw, K. (2000) A review of *Jatropha curcas*: an oil plant of unfulfilled promise. *Biomass and Bioenergy* 19:1-5.
- Onyema, M.C. Osuagwu, N.C. and Igbokwe, B.I. (2008) The development and use of biofuels as alternative energy sources: issues, prospects and challenges for Africa. In: Popoola, L., (ed.). Proceedings of the 32nd Annual Conference of the Forestry Association of Nigeria (FAN) held in Umuahia, Abia State, between 20th and 24th October, 2008. Ppp 675 – 681.
- Statistical software (SAS) (2005) Hargen and enhanced SAS Inst. Inc. USA.
- Sukarin, W., Yamada, Y and Sakaguchu, S. (1987) Characteristics of Physic nut, *Jatropha curcas* L.) as a new biomass crop in the tropics. *Japanese Agricultural Resources* 20(4): 302 – 303.
- Tigere, T.A., Gatsi, T.C., Mudita, I. I., Chikuvire, T.J., Thamangani, S. and Mavunganidze, Z. (2006) Potentials of *Jatropha curcas* in improving smallholder farmers' livelihoods in Zimbabwe: an explanatory study of Makosa ward, Mutoko District *Journal of Sustainable Development in Africa* 7:11-16.
- Vwioko, D. E., Anoliefo, G. O. and Fashemi, S. D. (2006) Metal concentration in plant tissues of *Ricinus communis* L (Castor oil) grown in soil contaminated with spent lubricating soil. *Journal of Applied Environmental Management* 10 (3): 127 – 134.
- Wini, S.P. Osman, M., Emmanuel, D.S. and Sreedevi, T.K. (2006) Improved livelihoods and environmental protection through biodiesel plantations in Asia. *Asian Biotechnology and Development Review* 8(2): 11-29.